

MULTIPLICATIVE FUNCTIONALS ON ENSEMBLES OF NON-INTERSECTING PATHS

ALEXEI BORODIN, IVAN CORWIN, AND DANIEL REMENIK

ABSTRACT. The purpose of this article is to develop a theory behind the occurrence of “path-integral” kernels in the study of extended determinantal point processes and non-intersecting line ensembles. Our first result shows how determinants involving such kernels arise naturally in studying ratios of partition functions and expectations of multiplicative functionals for ensembles of non-intersecting paths on weighted graphs. Our second result shows how Fredholm determinants with extended kernels (as arise in the study of extended determinantal point processes such as the Airy_2 process) are equal to Fredholm determinants with path-integral kernels. We also show how the second result applies to a number of examples including the stationary (GUE) Dyson Brownian motion, the Airy_2 process, the Pearcey process, the Airy_1 and $\text{Airy}_{2 \rightarrow 1}$ processes, and Markov processes on partitions related to the z -measures.

1. INTRODUCTION

The Airy_2 process is a universal scaling limit of a wide variety of probabilistic systems including random matrix theory, random growth processes, interacting particle systems and directed polymers in random media (see [24, 41] and references therein). Denoted $\text{Airy}_2(\cdot)$, it is defined via its consistent finite dimensional distributions: for $t_1 < t_2 < \dots < t_n$,

$$\mathbb{P}\left(\bigcap_{i=1}^n \{\text{Airy}_2(t_i) \leq s_i\}\right) = \det(I - \chi K_2^{\text{ext}})_{L^2(\{t_1, \dots, t_n\} \times \mathbb{R}, \mu)}. \quad (1)$$

Here χ is an operator which acts on functions $f: \{t_1, \dots, t_n\} \times \mathbb{R} \rightarrow \mathbb{R}$ as

$$\chi f(t_i, x) := \mathbf{1}_{x \geq s_i} f(t_i, x).$$

The operator K_2^{ext} acts as

$$K_2^{\text{ext}} f(t_i, x) := \sum_{j=1}^n \int_{\mathbb{R}} dy K_2^{\text{ext}}(t_i, x; t_j, y) f(t_j, y)$$

where $K_2^{\text{ext}}(s, x; t, y)$ is the “extended” Airy_2 kernel given by

$$K_2^{\text{ext}}(s, x; t, y) := \begin{cases} \int_0^\infty d\lambda e^{-\lambda(s-t)} \text{Ai}(x+\lambda) \text{Ai}(y+\lambda) & \text{if } s \geq t, \\ -\int_{-\infty}^0 d\lambda e^{-\lambda(s-t)} \text{Ai}(x+\lambda) \text{Ai}(y+\lambda) & \text{if } s < t, \end{cases}$$

with $\text{Ai}(x)$ the classical Airy function. The right-hand side of (1) is the Fredholm determinant of the identity minus a trace class operator (see Section 3.1 for definition and details) and the measure μ appearing there is the product of counting measure on $\{t_1, \dots, t_n\}$ and Lebesgue measure on \mathbb{R} .

The formula given in (1) for the finite dimensional distributions of the Airy_2 process becomes increasingly cumbersome as n increases. This is due to the n -dependence in the L^2 space on which the operators act. When taking a limit of a sequence of operators, or their determinants, it is convenient to have the operators all act on the same L^2 space, rather than a sequence of different spaces.

In Prähofer and Spohn's initial work on the Airy_2 process (see Section 5 of [36] for $n = 2$ or [22, 37, 40] for $n \geq 2$) the extended kernel formula is shown to be equivalent to the following “path-integral” kernel formula:

$$\mathbb{P}\left(\bigcap_{i=1}^n \{\text{Airy}_2(t_i) \leq s_i\}\right) = \det(I - K_2 + \bar{P}_{s_1} e^{(t_1-t_2)H} \bar{P}_{s_2} \cdots e^{(t_{n-1}-t_n)H} \bar{P}_{s_n} e^{(t_n-t_1)H} K_2)_{L^2(\mathbb{R})}. \quad (2)$$

Here $K_2(x, y) = K_2^{\text{ext}}(0, x; 0, y)$ is called the Airy_2 kernel, $\bar{P}_s g(x) = \mathbf{1}_{x \leq s} g(x)$ is a projection operator and $H = -\Delta + x$ is called the Airy Hamiltonian because $H\text{Ai}(\cdot - s) = s\text{Ai}(\cdot - s)$ (Δ is the Laplacian on \mathbb{R}). The dependence on n has been absorbed into the operator rather than the L^2 space, and it is now plausible to take a large n limit.

The reason we call this a path-integral kernel is because a portion of it can be written in terms of the expectation of a certain path-integral. By the Feynman-Kac formula (cf. [28]),

$$\bar{P}_{s_1} e^{(t_1-t_2)H} \bar{P}_{s_2} \cdots e^{(t_{n-1}-t_n)H} \bar{P}_{s_n} f(x) = \mathbb{E}_{b(t_1)=x} \left[f(b(t_n)) e^{-\int_{t_1}^{t_n} b(s) ds} \prod_{i=1}^n \mathbf{1}_{b(t_i) \leq s_i} \right], \quad (3)$$

where $b: [t_1, t_n] \rightarrow \mathbb{R}$ is the trajectory of a Brownian motion with diffusion coefficient 2 starting at $b(t_1) = x$.

Let $t_1 < \cdots < t_n$ fill out the interval $[\ell, r]$ and let $s_i = h(t_i)$ for some function $h: [\ell, r] \rightarrow \mathbb{R}$. Then as n goes to infinity, the above operator has a limit in trace norm (see [22] or Section 4.2 below for details) given by

$$\Gamma_{\ell, r}^h f(x) = \mathbb{E}_{b(\ell)=x} \left[f(b(r)) e^{-\int_{\ell}^r b(s) ds} \mathbf{1}_{b \leq h} \right],$$

where $\{b \leq h\}$ denotes the event $\{b(s) \leq h(s), \forall s \in [\ell, r]\}$. Thus, it is shown in [22] Theorem 2 or equation (41) below that

$$\mathbb{P}(\text{Airy}_2(s) \leq h(s), \forall s \in [\ell, r]) = \det(I - K_2 + \Gamma_{\ell, r}^h e^{(r-\ell)H} K_2)_{L^2(\mathbb{R})}. \quad (4)$$

The above formula proved useful in [22] in providing a direct proof that the value of the maximum of the Airy_2 process minus a parabola is distributed according to the (GOE) Tracy-Widom distribution; and in [32] in computing the joint distribution for the value and location (in t) of the maximum. The two-time path-integral kernel formula in [36] was utilized to compute asymptotics of the two-time covariance of the Airy_2 process, since the extended kernel does not easily yield this. Note that the left-hand side in the last formula presupposes the existence of a continuous version of the Airy_2 process. This was first shown to exist in [26].

What is remarkable about formula (4) is that the right-hand side is simple (despite the cumbersome finite dimensional distributions given above) and the event in question in the left-hand side has a clear translation into the operator $\Gamma_{\ell, r}^h$. As a further application of the Feynman-Kac formula as well as the Cameron-Martin-Girsanov formula (see [22]), the integral kernel of $\Gamma_{\ell, r}^h$ can be expressed as

$$\Gamma_{\ell, r}^h(x, y) = \mathbb{P}_{b(\ell)=x-\ell^2, b(r)=y-r^2} (b(s) \leq h(s) - s^2 \text{ for all } s \in [\ell, r]),$$

where b is now a Brownian bridge run from $x - \ell^2$ at time ℓ to $y - r^2$ at time r (this means that $\Gamma_{\ell, r}^h f(x) = \int dy \Gamma_{\ell, r}^h(x, y) f(y)$). In other words, the probability that the Airy_2 process hits a function h can be expressed as the Fredholm determinant of an operator which is partly expressed by the probability that a Brownian bridge hits the same function (minus a parabola).

We will now see how these formulas for the Airy_2 process are part of a more general result.

1.1. Extended kernels and the path-integral kernel in a general setting. There are many other examples of extended determinantal point processes (some given in Section 4) and our aim is to find path-integral kernel formulas for these other processes. This may have further applications, although we do not address them here. For example, besides the previous work of [37], in [40] a path-integral kernel formula was discovered for the Airy_1 process and used to prove existence of a continuous version of the process and its Hölder regularity.

When the Airy_2 process was introduced, it arose as the top layer of the multi-layer Airy_2 process, which will be denoted $\{\text{Airy}_2(i; t) : i \in \mathbb{Z}_{\geq 1}, t \in \mathbb{R}\}$ and is such that $\text{Airy}_2(i; t) > \text{Airy}_2(j; t)$ for $i < j$. As t varies, $\sum_{i=1}^{\infty} \delta_{\text{Airy}_2(i; t)}$ forms an \mathbb{R} -indexed collection of point processes which has the structure of an (extended) determinantal point process (see [7] and references therein) with correlation kernel $K_2^{\text{ext}}(s, x; t, y)$.

This has the following consequence. For $t_1 < \dots < t_n$ fix $q_{t_i} : \mathbb{R} \rightarrow \mathbb{R}$ and let $\bar{q}_{t_i} = 1 - q_{t_i}$. For $g : \{t_1, \dots, t_n\} \rightarrow \mathbb{R}$ define $q(g) := \prod_{i=1}^n q_{t_i}(g(t_i))$ and likewise define $\bar{q}(g)$. Then (given some conditions on q to ensure convergence – see Section 4.2 below)

$$\mathbb{E} \left[\prod_{i=1}^{\infty} \bar{q}(\text{Airy}_2(i; \cdot)) \right] = \det(I - QK_2^{\text{ext}})_{L^2(\{t_1, \dots, t_n\} \times \mathbb{R}, \mu)} \quad (5)$$

where $Qf(t_i, x) := q_{t_i}(x)f(t_i, x)$. The left-hand side above is referred to here as the expectation of a “multiplicative functional” of the multi-layer process. When $q_{t_i}(x) = \mathbf{1}_{x > s_i}$ (and hence $\bar{q}_{t_i}(x) = \mathbf{1}_{x \leq s_i}$), the above formula reduces to (1) with $Q = \chi$.

Our first result shows that the type of identity between extended and path-integral kernel Fredholm determinants which one gets by equating the right-hand sides of (1) and (2) is quite general and dependent on a few structural properties of the kernels.

In stating our results presently, we leave out a number of technical assumptions (see Section 3 for these details) and assume that we have the following collection of operators on functions $f : \{t_1, \dots, t_n\} \rightarrow \mathbb{R}$:

- For each $1 \leq i, j \leq n$, \mathcal{W}_{t_i, t_j} (with the convention $\mathcal{W}_{t_i, t_i} = I$);
- For each $1 \leq i \leq n$, K_{t_i} ;
- A diagonal operator Q such that $Qf(t_i, \cdot) := Q_{t_i}f(t_i, \cdot)$ where for $1 \leq i \leq n$ and $g : \mathbb{R} \rightarrow \mathbb{R}$, $Q_{t_i}g(x) := q_{t_i}(x)g(x)$.

Theorem 1.1 (Theorem 3.3, with technical assumptions suppressed). *Assume that for all $1 \leq i \leq j \leq k \leq n$ the following holds:*

- *Right-invertibility:* $\mathcal{W}_{t_i, t_j} \mathcal{W}_{t_j, t_i} K_{t_i} = K_{t_i}$;
- *Semigroup property:* $\mathcal{W}_{t_i, t_j} \mathcal{W}_{t_j, t_k} = \mathcal{W}_{t_i, t_k}$;
- *Reversibility relation:* $\mathcal{W}_{t_i, t_j} K_{t_j} = K_{t_i} \mathcal{W}_{t_i, t_j}$.

Then

$$\det(I - QK^{\text{ext}})_{L^2(\{t_1, \dots, t_n\} \times \mathbb{R}, \mu)} = \det(I - K_{t_1} + \bar{Q}_{t_1} \mathcal{W}_{t_1, t_2} \bar{Q}_{t_2} \cdots \mathcal{W}_{t_{n-1}, t_n} \bar{Q}_{t_n} \mathcal{W}_{t_n, t_1} K_{t_1})_{L^2(\mathbb{R})},$$

where

$$K^{\text{ext}}(t_i, x; t_j, y) = \begin{cases} \mathcal{W}_{t_i, t_j} K_{t_j}(x, y) & \text{if } i \geq j, \\ -\mathcal{W}_{t_i, t_j} (I - K_{t_j})(x, y) & \text{if } i < j, \end{cases}$$

and $\bar{Q}_{t_i} = I - Q_{t_i}$.

This is proved in Section 4.2 essentially via linear algebra.

Letting $\mathcal{W}_{t_i, t_j} = e^{-(t_j - t_i)H}$, $K_{t_i} = K_2$ and $Q_{s_i} = P_{s_i}$ we recover the equality (2) for the Airy_2 process. More generally, the result implies that the expectation of a multiplicative functional of the multi-layer Airy_2 process (5) can be expressed in a similar way, by replacing each P_{s_i} by Q_{s_i} on the right-hand side of (2).

In Section 4 we apply this theorem to a variety of examples of extended determinantal point processes such as the stationary (GUE) Dyson Brownian motion, the Airy_2 process, the Pearcey process, and Markov processes on partitions related to the z -measures. We also show how the identity applies to signed extended determinantal point processes such as the Airy_1 and $\text{Airy}_{2 \rightarrow 1}$ processes. In the case of the stationary (GUE) Dyson Brownian motion and the Airy_2 process, we also obtain the continuum limits of the corresponding path-integral kernel formulas (which is likely doable in other cases as well).

1.2. Ensembles of non-intersecting paths and the path-integral kernel. The multi-layer Airy_2 process arises as the scaling limit of a variety of ensembles of non-intersecting paths (see for instance [27]). The occurrence of extended kernel determinants in such ensembles is a consequence of the Eynard-Mehta theorem, which implies the existence of an extended determinantal point process structure [17, 23, 26, 33, 45]. The equivalence of the extended kernel determinant formula with the path-integral kernel determinant formula which is given in Theorem 1.1 (see also Theorem 3.3) is via linear algebra, but does not indicate why such a path-integral kernel formula exists. Theorem 1.3 below provides a direct link between ensembles of non-intersecting paths and path-integral kernel determinant formulas, and its proof boils down to the Lindström-Gessel-Viennot Lemma (recorded below as Lemma 2.1). By first proving the path-integral kernel determinant formula and then relating it to an extended kernel determinant formula, this provides another proof of the extended determinant point process structure for these ensembles (i.e., the Eynard-Mehta theorem), see Section 1.2.1.

Let us now introduce the ensembles of non-intersecting paths. Fix $T \in \mathbb{Z}_{\geq 0}$ and $N \in \mathbb{Z}_{\geq 1}$. Let $G = (V, E)$ be a finite directed acyclic planar graph with vertex set $V = V_0 \sqcup V_1 \sqcup \dots \sqcup V_T$ (here \sqcup represents the disjoint union of sets) and directed edge set $E = E_{0 \rightarrow 1} \sqcup E_{1 \rightarrow 2} \sqcup \dots \sqcup E_{T-1 \rightarrow T}$ where $E_{n \rightarrow n+1}$ only contains edges from $x \rightarrow y$ with $x \in V_n$ and $y \in V_{n+1}$. Here $x \rightarrow y$ denotes an edge directed from x to y .

As an example, let V_n be vertices of \mathbb{Z}^2 of the form (n, i) for $n-i \equiv 0 \pmod{2}$ (and $|i| \leq M$ for some M) and let E_n contain all directed edges from (x, n) to $(x \pm 1, n+1)$. Paths in this directed graph are trajectories of simple symmetric random walks (constrained to stay within distance M from the origin), cf. Figure 1.

We define a *path* π as a sequence of edges ($e_0 = x_0 \rightarrow x_1, e_1 = x_1 \rightarrow x_2, \dots, e_{T-1} = x_{T-1} \rightarrow x_T$) where $x_i \in V_i$ for $0 \leq i \leq T$. For such a path, let $\pi(n) = x_n$ denote the n^{th} vertex in the path.

To edges $e \in E$ we associate weights $w_e \in \mathbb{R}$, and to a path π we associate a weight $w(\pi)$ given by the product of the weights $w(e_n)$ along the edges e_n of π . For $x \in V_0$ and $y \in V_T$ we define a transition matrix

$$\mathcal{W}(x, y) := \sum_{\pi: x \rightarrow y} w(\pi)$$

where the summation is over all paths π from x to y . Instead of considering just a single path, we may consider ensembles of N non-intersecting paths from elements of V_0 to elements of V_T (by which we mean paths which use disjoint collections of vertices). We define the collection of all such paths as

$$\mathcal{NI}(N) := \left\{ \Pi = \{\pi_1, \dots, \pi_N\} : \forall i, \pi_i \text{ goes from } V_0 \text{ to } V_T \text{ and no two paths intersect} \right\}.$$

We will describe a measure on such an ensemble. This requires the introduction of two additional families of functions. For N fixed, consider functions $\psi_i: V_0 \rightarrow \mathbb{R}$, $1 \leq i \leq N$, and $\varphi_j: V_T \rightarrow \mathbb{R}$, $1 \leq j \leq N$. Define the weight of $\Pi \in \mathcal{N.I.}(N)$ as

$$Wt(\Pi) := \det[\psi_i(\pi_j(0))]_{i,j=1}^N \left(\prod_{i=1}^N w(\pi_i) \right) \det[\varphi_i(\pi_j(T))]_{i,j=1}^N, \quad (6)$$

and the partition function as

$$Z = \sum_{\Pi \in \mathcal{N.I.}(N)} Wt(\Pi).$$

If ψ_i and φ_j are δ -functions then the ends of the paths are fixed.

Assuming that $Z \neq 0$ we may define a measure (not necessarily positive but with total integral 1) on $\Pi \in \mathcal{N.I.}(N)$ as

$$\nu(\Pi) = \frac{Wt(\Pi)}{Z}.$$

When each of the three factors on the right-hand side of (6) are positive (and thus, in particular, ν is a probability measure), one may think of ν as follows. The determinants $\det[\psi_i(\pi_j(0))]_{i,j=1}^N$ and $\det[\varphi_i(\pi_j(T))]_{i,j=1}^N$ define measures on the collections of N initial points in V_0 and N final points in V_T . The weights $w(\pi_i)$ in the middle factor in (6) describe the transition probabilities for N independent paths (π_1, \dots, π_N) connecting these points. The measure ν is restricted to non-intersecting paths, and the division by the normalizing constant Z means that ν corresponds to a measure conditioned on the N paths not intersecting.

Define $\varphi_j^{(0)}: V_0 \rightarrow \mathbb{R}$ by $\varphi_j^{(0)}(x) := \sum_{y \in V_T} \mathcal{W}(x, y) \varphi_j(y)$. Note that this implies that \mathcal{W} has a right-inverse on $\text{span}\{\varphi_i^{(0)}\}_{i=1}^N$ which is given by $\mathcal{W}^{-1} \varphi_j^{(0)} = \varphi_j$.

We will make the following *biorthogonality assumption* on the $\{\psi_i\}_{i=1}^N$ and $\{\varphi_j^{(0)}\}_{j=1}^N$:

$$\sum_{x \in V_0} \psi_i(x) \varphi_j^{(0)}(x) = \mathbf{1}_{i=j}.$$

Remark 1.2. Assuming $Z \neq 0$, one can show that it is always possible to perform a linear transformation in $\text{span}\{\psi_i\}_{i=1}^N$ in such a way that the biorthogonality assumption is satisfied and ν remains unchanged.

The final concept we introduce is that of a *path-integral functional*, which is any function

$$f: E_{0 \rightarrow 1} \times E_{1 \rightarrow 2} \times \dots \times E_{T-1 \rightarrow T} \rightarrow \mathbb{R}$$

such that

$$f(e_0, e_1, \dots, e_{T-1}) = \prod_{n=0}^{T-1} f_n(e_n)$$

for functions $f_n: E_{n \rightarrow n+1} \rightarrow \mathbb{R}$. This definition extends to a directed path π from V_0 to V_T by setting $f(\pi)$ equal to f applied to the ordered sequence of edges in π . From the function f define a second set of edge weights $\tilde{w}_e := f_n(e) w_e$ where n is such that $e \in E_{n \rightarrow n+1}$. With respect to these weights $\{\tilde{w}_e\}_{e \in E}$ define a transition matrix

$$\widetilde{\mathcal{W}}(x, y) = \sum_{\pi: x \rightarrow y} \tilde{w}(\pi).$$

The following theorem is a consequence of Theorem 2.2, which is a similar result for a more general graph setting. The theorem shows how the path-integral kernel determinant naturally arises from ensembles of non-intersecting paths. A proof of the below result appears in Section 2.

Theorem 1.3. *For any path-integral functional f as above*

$$\sum_{\Pi=(\pi_1, \dots, \pi_N) \in \mathcal{NI}(N)} \prod_{i=1}^N f(\pi_i) \nu(\Pi) = \det(I - K + \widetilde{\mathcal{W}} \mathcal{W}^{-1} K)_{L^2(V_0)}$$

where $K: L^2(V_0) \rightarrow L^2(V_0)$ is given by its kernel

$$K(x_1, x_2) = \sum_{i=1}^N \varphi_i^{(0)}(x_1) \psi_i(x_2).$$

As we will explain in the proof of Corollary 1.4 below, the above result can also be seen as a consequence of Theorem 1.1 and the known determinantal structure for ensembles of non-intersecting paths. Instead, we provide a direct and simple linear algebraic proof of Theorem 1.3 using only the Lindström-Gessel-Viennot Lemma. This provides an explanation of the appearance of path-integral kernel formulas.

1.2.1. Recovering the determinantal structure. As an application of Theorems 1.1 and 1.3, let us see how to recover the determinantal structure of the ensemble of non-intersecting paths distributed according to the law ν . We would like to show that for any collection of vertices $\{x_1, \dots, x_k\} \in V$, the ν -measure of the set

$$\left\{ \Pi \in \mathcal{NI}(N) : \text{all of the } x_i \text{ are visited by paths in } \Pi \right\}$$

can be written as $\det[K(x_i, x_j)]_{i,j=1}^k$ for some fixed matrix K with rows and columns indexed by the set of vertices V . This property can be seen as a consequence of Corollary 1.4 below which we show.

Consider any collection of functions $q_n: V_n \rightarrow \mathbb{R}$, $0 \leq n \leq T-1$. Consider the space of matrices with rows and columns indexed by V , and for notational convenience denote $x \in V_n$ as (n, x) so that matrix elements of a matrix M are written as $M(n, x; m, y)$. Define a matrix Q so that $Qf(n, x) = q_n(x)f(n, x)$. For $m \leq n$ and $x \in V_m$, $y \in V_n$ define $\mathcal{W}_{m,n}(x, y) := \sum_{\pi: x \rightarrow y} w(\pi)$ (for $m = n$ let this be the identity matrix). For $x \in V_n$ define $\varphi_j^{(n)}(x) := \sum_{y \in V_T} \mathcal{W}_{n,T}(x, y) \varphi_j(y)$. For $x_1, x_2 \in V_n$ define $K_n(x_1, x_2) := \sum_{i=1}^N \varphi_i^{(n)}(x_1) \psi_i(x_2)$. Note that for $m \leq n$, $\mathcal{W}_{m,n}$ has a right-inverse on $\text{span}\{\varphi_i^{(m)}\}_{i=1}^N$ which is given by $\mathcal{W}_{m,n}^{-1} \varphi_j^{(m)} = \varphi_j^{(n)}$. We will write this inverse as $\mathcal{W}_{n,m}$. On account of this we may define the following (extended kernel) matrix

$$K^{\text{ext}}(m, x; n, y) = \begin{cases} \mathcal{W}_{m,n} K_n(x, y) & \text{if } m \geq n, \\ -\mathcal{W}_{m,n} (I - K_n)(x, y) & \text{if } m < n. \end{cases}$$

Corollary 1.4. *For any collection of functions $q_n: V_n \rightarrow \mathbb{R}$, $0 \leq n \leq T-1$*

$$\sum_{\Pi=(\pi_1, \dots, \pi_N) \in \mathcal{NI}(N)} \prod_{i=1}^N \prod_{n=0}^{T-1} \bar{q}_n(\pi_i(n)) \nu(\Pi) = \det(I - Q K^{\text{ext}})_{L^2(V)}. \quad (7)$$

where we recall that $\pi(n)$ denotes the vertex in V_n through which π passes and that $\bar{q}(x) = 1 - q(x)$.

This result is essentially a version of the Eynard-Mehta Theorem.

Proof. We use Theorem 3.3 (the more general version of Theorem 1.1). The technical assumptions are immediately satisfied since we are dealing with a finite vector space. The right-invertibility, semigroup property and reversibility relation are all readily checked from

the definitions of the $\mathcal{W}_{m,n}$ and K_n . As a consequence of that theorem we find that we may rewrite the right-hand side of (7) as

$$\text{RHS}(7) = \det(I - K_0 + \overline{Q}_0 \mathcal{W}_{0,1} \overline{Q}_1 \mathcal{W}_{1,2} \cdots \overline{Q}_{T-1} \mathcal{W}_{T-1,T} \mathcal{W}_{T,0} K_0)_{L^2(V_0)}, \quad (8)$$

where for $x \in V_n$, $\overline{Q}_n f(n, x) := \bar{q}_n(x) f(n, x)$.

Define a path-integral functional f so that for an edge $e \in E_{n \rightarrow n+1}$ from $x \in E_n$ to $y \in E_{n+1}$, $f_n(e) = \bar{q}_n(x)$. As a consequence, for any path π from V_0 to V_T , $f(\pi) = \prod_{n=0}^{T-1} \bar{q}_n(\pi(n))$. This observation and Theorem 1.3 then imply that we may rewrite the left-hand side of (7) as

$$\text{LHS}(7) = \det(I - K_0 + \widetilde{\mathcal{W}}_{0,T} \mathcal{W}_{T,0} K_0)_{L^2(V_0)}. \quad (9)$$

Note that we have introduced the subscripts on the right-hand side to be consistent with the notation introduced before the statement of the corollary. Due to the specific type of path-integral functional f , it is now straight-forward to see that

$$\widetilde{\mathcal{W}}_{0,T} = \overline{Q}_0 \mathcal{W}_{0,1} \overline{Q}_1 \mathcal{W}_{1,2} \cdots \overline{Q}_{T-1} \mathcal{W}_{T-1,T}.$$

This implies that the right-hand sides of equations (8) and (9) match and therefore completes the proof of the corollary. \square

1.3. Outline. In Section 2 we prove a result about ensembles of non-intersecting paths on directed graphs (which implies Theorem 1.3 above). In Section 3 we prove a general result (which implies Theorem 1.1 above) showing the equality between certain extended kernel and path-integral Fredholm determinants. In Section 4 we apply this equality between Fredholm determinants to a variety of extended kernels from the literature, and in Appendix A we check the technical assumptions necessary in order to do this.

1.4. Acknowledgements. We thank Senya Shlosman for interest in this project early on. AB was partially supported by the NSF grant DMS-1056390. IC was partially supported by the NSF grant DMS-1208998 as well as by the Clay Research Fellowship and by Microsoft Research through the Schramm Memorial Fellowship. DR was partially supported by the Natural Science and Engineering Research Council of Canada and by a Fields-Ontario Postdoctoral Fellowship, as well as by Fondecyt Grant 1120309 and Conicyt Basal-CMM project. DR is appreciative for MIT's hospitality during the visit in which this project was initiated.

2. NON-INTERSECTING DIRECTED PATHS ON WEIGHTED GRAPHS

2.1. A general combinatorial result. Let $G = (V, E)$ be a finite directed acyclic planar graph with vertices V and edges E . Fix source vertices $X = \{x_1, \dots, x_N\}$ and sink vertices $Y = \{y_1, \dots, y_N\}$. Fix edge weights w_e for each directed edge $e \in E$.

A directed path π is a sequence of vertices connecting a vertex in X to a vertex in Y via directed edges in E . Denote the source vertex of π by $\pi(b)$ and the sink vertex of π by $\pi(d)$ ¹. We say that two paths *intersect* if their vertex sets have non-empty intersection. To a directed path π we associate a weight $w(\pi)$ which is given by the product of w_e over edges e of π . Define

$$\mathcal{W}(x, y) = \sum_{\pi: x \rightarrow y} w(\pi) \quad (10)$$

where the sum is over directed paths π from source vertex x to sink vertex y .

¹We use b to denote the source, or base vertex; and d to denote the sink, or destination vertex.




FIGURE 1. An example of a graph G with source vertices \mathcal{X} on the left-hand side and sink vertices \mathcal{Y} on the right-hand side. Here there are $N = 3$ non-intersecting paths which are shown in grey, with starting points $\pi_i(b)$ and ending points $\pi_i(d)$ for $i = 1, 2, 3$.

Define the ensemble of N directed non-intersecting paths from the elements of X to the elements of Y as

$$\mathcal{N.I.}(N; X \rightarrow Y) = \{ \{ \pi_1, \dots, \pi_N \} : \forall i, \pi_i(b) \in X, \pi_i(d) \in Y \text{ and no two paths intersect} \}.$$

Write $\Pi = \{ \pi_1, \dots, \pi_N \}$ for an element of $\mathcal{N.I.}(N; X \rightarrow Y)$. Note that the non-intersection condition and the fact that X and Y have N elements ensures that $\{ \pi_1(b), \dots, \pi_N(b) \} = X$ and $\{ \pi_1(d), \dots, \pi_N(d) \} = Y$.

Lemma 2.1 ([25, 29, 31, 44]). *Fix $N \geq 1$. For any finite directed acyclic planar graph G with source vertices $X = \{x_1, \dots, x_N\}$, sink vertices $Y = \{y_1, \dots, y_N\}$ and edge weights w_e for each directed edge $e \in E$,*

$$\det[\mathcal{W}(x_i, y_j)]_{i,j=1}^N = \sum_{\Pi \in \mathcal{N.I.}(N; X \rightarrow Y)} \prod_{i=1}^N w(\pi_i).$$

Consider now finite sets of source vertices $\mathcal{X} \subset V$ and sink vertices $\mathcal{Y} \subset V$ (with at least N vertices in each of \mathcal{X} and \mathcal{Y}). For such \mathcal{X} and \mathcal{Y} we can likewise define the ensemble of N directed non-intersecting paths from elements of \mathcal{X} to elements of \mathcal{Y} . Denote this ensemble $\mathcal{N.I.}(N; \mathcal{X} \rightarrow \mathcal{Y})$.

Fix functions $\psi_i : \mathcal{X} \rightarrow \mathbb{R}$ for $1 \leq i \leq N$ and functions $\varphi_j : \mathcal{Y} \rightarrow \mathbb{R}$ for $1 \leq j \leq N$. For $\Pi = \{ \pi_1, \dots, \pi_N \}$ with source vertices $\{ \pi_1(b), \dots, \pi_N(b) \} \subset \mathcal{X}$ and sink vertices $\{ \pi_1(d), \dots, \pi_N(d) \} \subset \mathcal{Y}$ we define the weight of Π as

$$Wt(\Pi) := \det[\psi_i(\pi_j(b))]_{i,j=1}^N \left(\prod_{i=1}^N w(\pi_i) \right) \det[\varphi_i(\pi_j(d))]_{i,j=1}^N.$$

Define a partition function for directed non-intersecting ensembles of N paths from \mathcal{X} to \mathcal{Y} with respect to weights $\{w_e\}_{e \in E}$ and functions $\{\psi_i\}_{i=1}^N, \{\varphi_j\}_{j=1}^N$ as

$$Z = Z(\mathcal{X}, \mathcal{Y}, \{w_e\}_{e \in E}, \{\psi_i\}_{i=1}^N, \{\varphi_j\}_{j=1}^N) := \sum_{\Pi \in \mathcal{N.I.}(N; \mathcal{X} \rightarrow \mathcal{Y})} Wt(\Pi).$$

For $1 \leq j \leq N$ define $\varphi_j^{(b)} : \mathcal{X} \rightarrow \mathbb{R}$ by

$$\varphi_j^{(b)}(x) := \sum_{y \in \mathcal{Y}} \mathcal{W}(x, y) \varphi_j(y) \quad (11)$$

and further define the operator $K : L^2(\mathcal{X}) \rightarrow L^2(\mathcal{X})$ by its kernel (which is just an \mathcal{X} -indexed matrix)

$$K(x_1, x_2) = \sum_{i=1}^N \varphi_i^{(b)}(x_1) \psi_i(x_2). \quad (12)$$

Observe that \mathcal{W} has a right-inverse on $\text{span}\{\varphi_i^{(b)}\}_{i=1}^N$, which we will denote by \mathcal{W}^{-1} , given by

$$\mathcal{W}^{-1} \varphi_j^{(b)} = \varphi_j.$$

In particular, since the range of K is contained in $\text{span}\{\varphi_i^{(b)}\}_{i=1}^N$, $\mathcal{W}^{-1}K$ is well defined as an operator mapping $L^2(\mathcal{X})$ to $L^2(\mathcal{Y})$:

$$\mathcal{W}^{-1}Kf = \sum_{i=1}^N \langle \psi_i, f \rangle_{L^2(\mathcal{X})} \varphi_i, \quad (13)$$

where $\langle \cdot, \cdot \rangle_{L^2(\mathcal{X})}$ is the inner product in $L^2(\mathcal{X})$.

We say that the *biorthogonality assumption* is satisfied if

$$\left\langle \psi_i, \varphi_j^{(b)} \right\rangle_{L^2(\mathcal{X})} = \mathbf{1}_{i=j} \quad \text{for all } 1 \leq i, j \leq N.$$

Theorem 2.2. *Let $G = (V, E)$ be a finite directed acyclic planar graph. Fix sets of source vertices $\mathcal{X} \subset V$ and sink vertices $\mathcal{Y} \subset V$. Fix edge weights w_e and a second set of weights \tilde{w}_e for each directed edge $e \in E$. Fix functions $\psi_i : \mathcal{X} \rightarrow \mathbb{R}$ for $1 \leq i \leq N$ and functions $\varphi_j : \mathcal{Y} \rightarrow \mathbb{R}$ for $1 \leq j \leq N$ which satisfy the biorthogonality assumption with $\varphi_j^{(b)}$ defined via the w_e weights. Write*

$$Z = Z(\mathcal{X}, \mathcal{Y}, \{w_e\}_{e \in E}, \{\psi_i\}_{i=1}^N, \{\varphi_j\}_{j=1}^N) \quad \text{and} \quad \tilde{Z} = Z(\mathcal{X}, \mathcal{Y}, \{\tilde{w}_e\}_{e \in E}, \{\psi_i\}_{i=1}^N, \{\varphi_j\}_{j=1}^N).$$

Then

$$\frac{\tilde{Z}}{Z} = \det(I - K + \widetilde{\mathcal{W}}\mathcal{W}^{-1}K)_{L^2(\mathcal{X})},$$

where $\widetilde{\mathcal{W}} : L^2(\mathcal{Y}) \rightarrow L^2(\mathcal{X})$ is given by (10) with w replaced by \tilde{w} , and $\mathcal{W}^{-1}K : L^2(\mathcal{X}) \rightarrow L^2(\mathcal{Y})$ is defined in (13).

Remark 2.3. As will be clear from the proof, the biorthogonality assumption implies that $Z \neq 0$. Conversely, one can show that if $Z \neq 0$ then there exists a linear change of basis in the space spanned by $\{\psi_i\}_{i=1}^N$ (or equally well the space spanned by $\{\varphi_j\}_{j=1}^N$) which leads back to the biorthogonality assumption being satisfied and does not change the ratio \tilde{Z}/Z .

Before turning to the proof of the theorem, let us check that it implies Theorem 1.3.

Proof of Theorem 1.3. Recall that for the path-integral functional f , we defined a set of weights $\tilde{w}_e = f_n(e)w_e$ where $e \in E_{n \rightarrow n+1}$. Let $Z = Z(V_0, V_T, \{w_e\}_{e \in E}, \{\psi_i\}_{i=1}^N, \{\varphi_j\}_{j=1}^N)$ and $\tilde{Z} = Z(V_0, V_T, \{\tilde{w}_e\}_{e \in E}, \{\psi_i\}_{i=1}^N, \{\varphi_j\}_{j=1}^N)$. We claim that

$$\int_{\Pi \in \mathcal{N}(\mathcal{I}(N; V_0 \rightarrow V_T))} d\nu(\Pi) \prod_{i=1}^N f(\pi_i) = \frac{\tilde{Z}}{Z} = \det(I - K + \widetilde{\mathcal{W}}\mathcal{W}^{-1}K)_{L^2(V_0)}.$$

The second equality is an immediate corollary of Theorem 2.2. To see the first equality above observe that

$$\begin{aligned} \frac{\tilde{Z}}{Z} &= \sum_{\Pi \in \mathcal{N.I.}(N; V_0 \rightarrow V_T)} \prod_{i=1}^N \prod_{n=0}^{T-1} f_n(\pi_i(n) \rightarrow \pi_i(n+1)) \frac{Wt(\Pi)}{Z} \\ &= \int_{\Pi \in \mathcal{N.I.}(N; V_0 \rightarrow V_T)} d\nu(\Pi) \prod_{i=1}^N \prod_{n=0}^{T-1} f_n(\pi_i(n) \rightarrow \pi_i(n+1)) = \int_{\Pi \in \mathcal{N.I.}(N; V_0 \rightarrow V_T)} d\nu(\Pi) \prod_{i=1}^N f(\pi_i) \end{aligned}$$

as desired. \square

Proof of Theorem 2.2. The proof is linear algebra. We may rewrite \tilde{Z} by first summing over the subsets of \mathcal{X} and \mathcal{Y} which host the source and sink vertices, and then considering all non-intersecting paths between these sets. Thus

$$\begin{aligned} \tilde{Z} &= \sum_{\substack{X=\{x_1, x_2, \dots, x_N\} \subset \mathcal{X} \\ Y=\{y_1, y_2, \dots, y_N\} \subset \mathcal{Y}}} \det[\psi_i(x_j)]_{i,j=1}^N \left(\sum_{\Pi \in \mathcal{N.I.}(N; X \rightarrow Y)} \prod_{i=1}^N \tilde{w}(\pi_i) \right) \det[\varphi_i(y_j)]_{i,j=1}^N \\ &= \sum_{\substack{\{x_1, x_2, \dots, x_N\} \subset \mathcal{X} \\ \{y_1, y_2, \dots, y_N\} \subset \mathcal{Y}}} \det[\psi_i(x_j)]_{i,j=1}^N \left(N! \det[\tilde{\mathcal{W}}(x_i, y_j)]_{i,j=1}^N \right) \det[\varphi_i(y_j)]_{i,j=1}^N. \end{aligned}$$

The second line follows by an application of Lemma 2.1.

We may now apply the Cauchy-Binet identity twice. The first application is with respect to the summation in the x 's, and it yields

$$\sum_{\{x_1, x_2, \dots, x_N\} \subset \mathcal{X}} \det[\psi_i(x_j)]_{i,j=1}^N \det[\tilde{\mathcal{W}}(x_i, y_j)]_{i,j=1}^N = \det \left[\sum_{x \in \mathcal{X}} \psi_i(x) \tilde{\mathcal{W}}(x, y_j) \right]_{i,j=1}^N.$$

The second application likewise is applied to the summation in y 's and yields that

$$\tilde{Z} = \det \left[\sum_{\substack{x \in \mathcal{X} \\ y \in \mathcal{Y}}} \psi_i(x) \tilde{\mathcal{W}}(x, y) \varphi_j(y) \right]_{i,j=1}^N. \quad (14)$$

Observe that by the same argument we can obtain an analogous expression for Z with $\tilde{\mathcal{W}}$ replaced by \mathcal{W} . However, by the definition of $\varphi_j^{(b)}$ we find that

$$Z = \det \left[\sum_{x \in \mathcal{X}} \psi_i(x) \varphi_j^{(b)}(x) \right]_{i,j=1}^N.$$

By the biorthogonality assumption,

$$\sum_{x \in \mathcal{X}} \psi_i(x) \varphi_j^{(b)}(y) = \langle \psi_i, \varphi_j^{(b)} \rangle_{L^2(\mathcal{X})} = \mathbf{1}_{i=j},$$

and hence $Z = 1$. Now observe that we can rewrite the kernel in appearing in (14) as

$$\sum_{\substack{x \in \mathcal{X} \\ y \in \mathcal{Y}}} \psi_i(x) \tilde{\mathcal{W}}(x, y) \varphi_j(y) = \langle \tilde{\mathcal{W}} \varphi_i, \psi_j \rangle_{L^2(\mathcal{X})}$$

by using the definition of the operator $\tilde{\mathcal{W}}$ and the inner product on $L^2(\mathcal{X})$.

Therefore, it remains to prove that

$$\det(I - K + \widetilde{\mathcal{W}}\mathcal{W}^{-1}K)_{L^2(\mathcal{X})} = \det \left[\left\langle \widetilde{\mathcal{W}}\varphi_i, \psi_j \right\rangle_{L^2(\mathcal{X})} \right]_{i,j=1}^N. \quad (15)$$

To prove the above statement we will write down the matrix for the operator $I - K + \widetilde{\mathcal{W}}\mathcal{W}^{-1}K$ in the basis $(\varphi_1^{(b)}, \dots, \varphi_N^{(b)}, (\text{span}\{\psi_i\}_{i=1}^N)^\perp)$, where $(\text{span}\{\psi_i\}_{i=1}^N)^\perp$ represents any basis of the orthogonal complement of $\text{span}\{\psi_i\}_{i=1}^N$ in $L^2(\mathcal{X})$. Let us consider the action of the operator K on the basis elements. On $\varphi_j^{(b)}$ one sees that K acts as the identity operator:

$$\begin{aligned} (K\varphi_j^{(b)})(x_1) &= \sum_{x_2 \in \mathcal{X}} \left(\sum_{i=1}^N \varphi_i^{(b)}(x_1) \psi_i(x_2) \right) \varphi_j^{(b)}(x_2) = \sum_{i=1}^N \varphi_i^{(b)}(x_1) \left(\sum_{x_2 \in \mathcal{X}} \psi_i(x_2) \varphi_j^{(b)}(x_2) \right) \\ &= \sum_{i=1}^N \varphi_i^{(b)}(x_1) \left\langle \psi_i, \varphi_j^{(b)} \right\rangle_{L^2(\mathcal{X})} = \sum_{i=1}^N \varphi_i^{(b)}(x_1) \mathbf{1}_{i=j} = \varphi_j^{(b)}(x_1). \end{aligned}$$

It is likewise clear that K acts on the basis elements of $(\text{span}\{\psi_i\}_{i=1}^N)^\perp$ by taking them all to zero. Thus we may write K as the matrix

$$K = \left(\begin{array}{c|c} I & 0 \\ \hline 0 & 0 \end{array} \right)$$

where the two blocks correspond to the basis elements $\{\varphi_j^{(b)}\}_{j=1}^N$ and $(\text{span}\{\psi_i\}_{i=1}^N)^\perp$. This shows that

$$I - K = \left(\begin{array}{c|c} 0 & 0 \\ \hline 0 & I \end{array} \right).$$

The remaining operator to study is $\widetilde{\mathcal{W}}\mathcal{W}^{-1}K$. Writing the corresponding matrix in blocks as above we get from (13) that

$$\widetilde{\mathcal{W}}\mathcal{W}^{-1}K = \left(\begin{array}{c|c} A & 0 \\ \hline * & 0 \end{array} \right)$$

where the $N \times N$ matrix A is yet to be determined. The value of the star is not important. To see this, write

$$I - K + \widetilde{\mathcal{W}}\mathcal{W}^{-1}K = \left(\begin{array}{c|c} A & 0 \\ \hline * & I \end{array} \right)$$

and observe then that

$$\det(I - K + \widetilde{\mathcal{W}}\mathcal{W}^{-1}K)_{L^2(\mathcal{X})} = \det[A_{i,j}]_{i,j=1}^N. \quad (16)$$

The value of $A_{i,j}$ can be found by using the inner product,

$$A_{i,j} = \left\langle \widetilde{\mathcal{W}}\mathcal{W}^{-1}K\varphi_i^{(b)}, \psi_j \right\rangle_{L^2(\mathcal{X})}.$$

Recalling that $\mathcal{W}^{-1}K\varphi_i^{(b)} = \varphi_i$ we deduce that

$$A_{i,j} = \left\langle \widetilde{\mathcal{W}}\varphi_i, \psi_j \right\rangle_{L^2(\mathcal{X})}.$$

Combining this with (16) proves (15) and hence completes the proof of the theorem. \square

3. EQUIVALENCE OF EXTENDED KERNEL AND PATH-INTEGRAL KERNEL FREDHOLM DETERMINANTS

There are various types of limits one can take of graph-based non-intersecting line ensembles. In this section we will show that formulas of the type given in the previous section survive these limits. We do not prove this directly via a limit transition, but rather show how such formulas arise via manipulations of the extended kernel Fredholm determinants which describe these limiting systems. The main result of this section is, therefore, the equality of two types of Fredholm determinants.

This equality will be stated in an abstract setting in this section, and later applied to the examples we are interested in in Section 4. A concrete example to keep in mind is the Airy_2 process, which we will use throughout this section to illustrate the objects we will introduce and the assumptions we will make on them. The Airy_2 process was introduced in the Introduction and is discussed in further detail in Section 4.2, we refer the reader there for details and just recall the definitions of the Airy Hamiltonian $H = -\Delta + x$ and the Airy kernel $K_2(x, y) = \int_0^\infty d\lambda \text{Ai}(x + \lambda) \text{Ai}(y + \lambda)$.

3.1. Fredholm determinants. Let us briefly introduce some of the basic notions related to Fredholm determinants (we refer the reader to [43] for more details). Consider a separable Hilbert space \mathcal{H} and let A be a bounded linear operator acting on \mathcal{H} ($\mathcal{H} = L^2(\mathbb{R})$ in the Airy_2 case). Let $|A| = \sqrt{A^*A}$ be the unique positive square root of the operator A^*A . The *trace norm* of A is defined as $\|A\|_1 = \sum_{n=1}^\infty \langle e_n, |A|e_n \rangle$, where $\{e_n\}_{n \geq 1}$ is any orthonormal basis of \mathcal{H} . We say that $A \in \mathcal{B}_1(\mathcal{H})$, the family of *trace class operators*, if $\|A\|_1 < \infty$. For $A \in \mathcal{B}_1(\mathcal{H})$, one can define the trace $\text{tr}(A) = \sum_{n=1}^\infty \langle e_n, Ae_n \rangle$. For later use we also define the *Hilbert-Schmidt norm* $\|A\|_2 = \sqrt{\text{tr}(|A|^2)}$ and say that $A \in \mathcal{B}_2(\mathcal{H})$, the family *Hilbert-Schmidt operators*, if $\|A\|_2 < \infty$. Given $A \in \mathcal{B}_1(\mathcal{H})$ one can define a generalization of the finite-dimensional determinant, the *Fredholm determinant* $\det(I + A)_\mathcal{H}$. We refer the reader to [43] for the details of the definition in this level of generality and just point out that, as expected, $\det(I + A)_\mathcal{H} = \prod_n (1 + \lambda_n)$, where λ_n are the eigenvalues of A (counted with algebraic multiplicity).

The result presented in this section (Theorem 3.3) can be stated, under some conditions, for operators acting on a general separable Hilbert space. Nevertheless, in order to keep the presentation as simple as possible, and since it is the setting we need for the examples in Section 4, we will restrict ourselves to the case of integral operators on an L^2 space.

More precisely, we assume we are given a measure space (X, Σ, μ) and consider the Hilbert space $L^2(X, \mu)$. For brevity we will drop μ from the notation. We will also denote by $\mathcal{M}(X)$ the space of real-valued measurable functions on X . By an *integral operator* we mean an operator $A: D \subseteq \mathcal{M}(X) \rightarrow \mathcal{M}(X)$ acting as $Af(x) = \int_X \mu(dy) A(x, y)f(y)$, where $A: X \times X \rightarrow \mathbb{R}$ is the *integral kernel* of A . We will often speak interchangeably of an integral operator and its kernel. In particular we have abused notation by using the same letter to denote an integral operator and its kernel. We recall that the product of two integral operators is defined by $AB(x, y) = \int_X \mu(dz) A(x, z)B(z, y)$.

Though we will not appeal to this, we note that the Fredholm determinant $\det(I - K)_{L^2(X)}$ of a trace class operator $K: L^2(X) \rightarrow L^2(X)$ with continuous (in both x and y) integral kernel $K(x, y)$ has the following (absolutely convergent) series expansion

$$\det(I - K)_{L^2(X)} = 1 + \sum_{k \geq 1} \frac{(-1)^k}{k!} \int_X d\mu(x_1) \cdots \int_X d\mu(x_k) \det[K(x_i, x_j)]_{i,j=1}^k. \quad (17)$$

3.2. Assumptions for the theorem. In order to state the main theorem of this section in a fairly broad context, we must introduce a few operators and impose certain assumptions

upon them. Most of the assumptions are technical and intended to ensure well-definedness or finiteness of the various quantities involved in the statement of the theorem. The main (not just technical) assumption is given in Assumption 2.

Fix $t_1 < \dots < t_n$ for the duration of this section. We will be interested in comparing the Fredholm determinant of certain integral operators acting on the Hilbert spaces $L^2(X)$ and $L^2(\{t_1, \dots, t_n\} \times X)$ (the measure we use in the second space is the product of the counting measure on $\{t_1, \dots, t_n\}$ and μ). The operators we consider will be constructed from the following four families of operators:

- For each $1 \leq i < j \leq n$, an integral kernel \mathcal{W}_{t_i, t_j} (for convenience we also introduce the notation $\mathcal{W}_{t_i, t_i} = I$);
- For each $1 \leq i \leq n$, an integral kernel K_{t_i} ;
- For each $1 \leq i < j \leq n$, an integral kernel $\mathcal{W}_{t_j, t_i} K_{t_i}$ (for convenience we also introduce the notation $\mathcal{W}_{t_i, t_i} K_{t_i} = K_{t_i}$);
- For each $1 \leq i \leq n$, a multiplication operator Q_{t_i} acting on $\mathcal{M}(X)$ as $Q_{t_i} f(x) = q_{t_i}(x) f(x)$ for some $q_{t_i} \in \mathcal{M}(X)$.

The reason for the choice of notation $\mathcal{W}_{t_j, t_i} K_{t_i}$ in the third family of operators is that we will assume below that $\mathcal{W}_{t_i, t_j} \mathcal{W}_{t_j, t_i} K_{t_i} = K_{t_i}$ for $i < j$ (so that even though it is not defined above as its own operator, \mathcal{W}_{t_j, t_i} can be thought of as a right inverse of \mathcal{W}_{t_i, t_j} on the range of K_{t_i}).

We make the following (technical) assumption.

Assumption 1.

- The integral operators $Q_{t_i} \mathcal{W}_{t_i, t_j}$, $Q_{t_i} K_{t_i}$, $Q_{t_i} \mathcal{W}_{t_i, t_j} K_{t_j}$ and $Q_{t_j} \mathcal{W}_{t_j, t_i} K_{t_i}$ for $1 \leq i < j \leq n$ are all bounded operators mapping $L^2(X)$ to itself.
- The operator

$$K_{t_1} - \overline{Q}_{t_1} \mathcal{W}_{t_1, t_2} \overline{Q}_{t_2} \cdots \mathcal{W}_{t_{n-1}, t_n} \overline{Q}_{t_n} \mathcal{W}_{t_n, t_1} K_{t_1},$$

where $\overline{Q}_{t_i} = I - Q_{t_i}$, is a bounded operator mapping $L^2(X)$ to itself.

The last operator in the assumption will appear in the formula provided in Theorem 3.3. An alternative expression for this operator, which is in some cases more convenient for checking the assumption, is given in Lemma 3.1.

In the case of the Airy₂ process we take $X = \mathbb{R}$, choose μ to be the Lebesgue measure and set $\mathcal{W}_{t_i, t_j} = e^{(t_i - t_j)H}$, $K_{t_i} = K_2$, and $\mathcal{W}_{t_j, t_i} K_{t_i} = e^{(t_j - t_i)H} K_2$ for $1 \leq i \leq j \leq n$. One can take for example the operators Q_{t_i} to be projections on intervals $[a_i, \infty)$, that is, $Q_{t_i} f(x) = \mathbf{1}_{x \geq a_i} f(x)$, which corresponds to studying the finite dimensional distributions of the Airy₂ process (we will make a more general choice in Section 4.2).

Going back to the general setting, we will make a certain algebraic assumption on the operators \mathcal{W}_{t_i, t_j} , K_{t_i} and $\mathcal{W}_{t_j, t_i} K_{t_i}$.

Assumption 2. For each $i \leq j \leq k$ the following hold:

- Right-invertibility:* $\mathcal{W}_{t_i, t_j} \mathcal{W}_{t_j, t_i} K_{t_i} = K_{t_i}$;
- Semigroup property:* $\mathcal{W}_{t_i, t_j} \mathcal{W}_{t_j, t_k} = \mathcal{W}_{t_i, t_k}$;
- Reversibility relation:* $\mathcal{W}_{t_i, t_j} K_{t_j} = K_{t_i} \mathcal{W}_{t_i, t_j}$.

The second property is clear in the Airy₂ case, while (i) and (iii) follow from the fact that K_2 is the projection operator into the negative (generalized) eigenspace of the Airy Hamiltonian H (see Section 4.2).

Let us now explain how these operators will be used. Using the kernels introduced above we define an *extended kernel* K^{ext} as follows: for $1 \leq i, j \leq n$ and $x, y \in X$,

$$K^{\text{ext}}(t_i, x; t_j, y) = \begin{cases} \mathcal{W}_{t_i, t_j} K_{t_j}(x, y) & \text{if } i \geq j, \\ -\mathcal{W}_{t_i, t_j} (I - K_{t_j})(x, y) & \text{if } i < j. \end{cases} \quad (18)$$

This definition coincides with the usual notion of extended correlation kernels of determinantal point processes, cf. [7, 17, 23, 26, 33, 45]. In the case of the Airy_2 process, it coincides with the definition given in the Introduction and in (40). As an operator, K^{ext} acts on $f \in L^1_{\text{loc}}(\{t_1, \dots, t_n\} \times X)$ as

$$K^{\text{ext}} f(t_i, x) = \sum_{j=1}^n \int_X d\mu(y) K^{\text{ext}}(t_i, x; t_j, y) f(t_j, y).$$

See Section 4 for concrete examples.

We also need to make the following (technical) analytical assumption.

Assumption 3. One can choose multiplication operators V_{t_i} , V'_{t_i} , U_{t_i} and U'_{t_i} acting on $\mathcal{M}(X)$, for $1 \leq i \leq n$, in such a way that:

- (i) $V'_{t_i} V_{t_i} Q_{t_i} = Q_{t_i}$ and $K_{t_i} U'_{t_i} U_{t_i} = K_{t_i}$, for all $1 \leq i \leq n$.
- (ii) The operators $V_{t_i} Q_{t_i} K_{t_i} V'_{t_i}$, $V_{t_i} Q_{t_i} \mathcal{W}_{t_i, t_j} V'_{t_j}$, $V_{t_i} Q_{t_i} \mathcal{W}_{t_i, t_j} K_{t_j} V'_{t_j}$ and $V_{t_j} Q_{t_j} \mathcal{W}_{t_j, t_i} K_{t_i} V'_{t_i}$ preserve $L^2(X)$ and are trace class in $L^2(X)$, for all $1 \leq i < j \leq n$.
- (iii) The operator $U_{t_i} [\mathcal{W}_{t_i, t_1} K_{t_1} - \overline{Q}_{t_i} \mathcal{W}_{t_i, t_{i+1}} \cdots \overline{Q}_{t_{n-1}} \mathcal{W}_{t_{n-1}, t_n} \overline{Q}_{t_n} \mathcal{W}_{t_n, t_1} K_{t_1}] U'_{t_1}$ preserves $L^2(X)$ and is trace class in $L^2(X)$, for all $1 \leq i \leq n$, where $\overline{Q}_{t_i} = I - Q_{t_i}$.

The primes in U'_{t_i} and V'_{t_i} mean that these are almost (left) inverses of the operators U_{t_i} and V_{t_i} , and hence the multiplication by these operators in (ii) and (iii) should be thought of as a conjugation. The distinction is because in many cases it will be necessary to let V_{t_i} be multiplication by a function which is 0 where q_{t_i} is 0, in which case V_{t_i} is not invertible, with an analogous situation for U_{t_i} and U'_{t_i} .

Before stating the main result of this section, Theorem 3.3, let us state a formula which reexpresses the operator appearing in Assumption 3(iii). Besides being used in the proof of the below theorem, this formula is often useful in checking the assumption (for example, as in Remark 3.2).

Lemma 3.1. *Writing $\overline{Q}_t = I - Q_t$, we have, for any $1 \leq i \leq n$,*

$$\begin{aligned} & \mathcal{W}_{t_i, t_1} K_{t_1} - \overline{Q}_{t_i} \mathcal{W}_{t_i, t_2} \overline{Q}_{t_2} \cdots \mathcal{W}_{t_{n-1}, t_n} \overline{Q}_{t_n} \mathcal{W}_{t_n, t_1} K_{t_1} \\ &= \sum_{j=i}^n \sum_{k=0}^{n-j} (-1)^k \sum_{j=a_0 < a_1 < \cdots < a_k \leq n} \mathcal{W}_{t_i, t_j} Q_{t_j} \mathcal{W}_{t_j, t_{a_1}} Q_{t_{a_1}} \mathcal{W}_{t_{a_1}, t_{a_2}} Q_{t_{a_2}} \cdots \mathcal{W}_{t_{a_{k-1}}, t_{a_k}} Q_{t_{a_k}} \mathcal{W}_{t_{a_k}, t_1} K_{t_1}. \end{aligned}$$

We postpone the proof of this lemma until the end of this section.

Remark 3.2. Suppose that there exist multiplication operators \tilde{V}_{t_i} and \tilde{V}'_{t_i} acting on $\mathcal{M}(X)$, for $1 \leq i \leq n$, in such a way that:

- (i) $\tilde{V}'_{t_i} \tilde{V}_{t_i} Q_{t_i} = Q_{t_i}$ and $K_{t_i} \tilde{V}_{t_i} \tilde{V}'_{t_i} = K_{t_i}$, for all $1 \leq i \leq n$;
- (ii) The operators $\tilde{V}_{t_i} Q_{t_i} K_{t_i} \tilde{V}'_{t_i}$, $\tilde{V}_{t_i} Q_{t_i} \mathcal{W}_{t_i, t_j} \tilde{V}'_{t_j}$, $\tilde{V}_{t_i} Q_{t_i} \mathcal{W}_{t_i, t_j} K_{t_j} \tilde{V}'_{t_j}$ and $\tilde{V}_{t_j} Q_{t_j} \mathcal{W}_{t_j, t_i} K_{t_i} \tilde{V}'_{t_i}$ preserve $L^2(X)$ and are trace class in $L^2(X)$, for all $1 \leq i < j \leq n$.

Then it is not hard to check, using the formula given in Lemma 3.1, that Assumption 3 holds, taking $U_{t_i} = V'_{t_i} = \tilde{V}'_{t_i}$ and $U'_{t_i} = V_{t_i} = \tilde{V}_{t_i}$ (see the end of the proof of Corollary 4.6

in Appendix A for more details). In the case of the Airy_2 process, when the operators Q_{t_i} are of the form $Q_{t_i}f(x) = \mathbf{1}_{x \geq a_i}f(x)$ as discussed above, both \tilde{V}_{t_i} and \tilde{V}'_{t_i} can be taken to be the identity. If, on the other hand, one assumes $q_{t_i}(x)$ to be 0 for $x < a_i$ but to grow at a certain rate for $x \geq a_i$, as we will in Section 4.2, then it is necessary to choose these operators more carefully (see the proof of Corollary 4.6).

3.3. Identity between extended and path-integral kernel Fredholm determinants. Define a diagonal operator Q acting on $f \in \mathcal{M}(\{t_1, \dots, t_n\} \times X)$ as

$$Qf(t_i, \cdot) = Q_{t_i}f(t_i, \cdot). \quad (19)$$

Note that, by Assumption 1, QK^{ext} preserves $L^2(\{t_1, \dots, t_n\} \times X)$. The following result expresses the Fredholm determinant of $I - QK^{\text{ext}}$ on $L^2(\{t_1, \dots, t_n\} \times X)$ as a Fredholm determinant on $L^2(X)$. The first example of such a formula was provided by [36] for the case of the Airy_2 process (see also [37]). This was later extended to the Airy_1 process in [40]. This type of formulas have recently been found to be very useful in the study of these processes, see for example [22, 32, 38–40].

Theorem 3.3. *With the above notation, and under Assumptions 1, 2 and 3, we have*

$$\det(I - QK^{\text{ext}})_{L^2(\{t_1, \dots, t_n\} \times X)} = \det(I - K_{t_1} + \overline{Q}_{t_1} \mathcal{W}_{t_1, t_2} \overline{Q}_{t_2} \cdots \mathcal{W}_{t_{n-1}, t_n} \overline{Q}_{t_n} \mathcal{W}_{t_n, t_1} K_{t_1})_{L^2(X)}, \quad (20)$$

where $\overline{Q}_t = I - Q_t$.

Remark 3.4. The operators appearing in both Fredholm determinants preserve $L^2(X)$ by Assumption 1. Moreover, the Fredholm determinants are well-defined thanks to Assumption 3, even though the operators appearing there are not necessarily trace class. In fact, if we define the diagonal operator V acting on $u \in L^2(\{t_1, \dots, t_n\} \times X)$ as $(Vu)_{t_i} = V_{t_i}u_{t_i}$, and similarly define V' , then $VQK^{\text{ext}}V'$ is trace class by Assumption 3(ii) and by the cyclic property of the determinant and the fact that $V'VQ = Q$ it leads to the same Fredholm expansion for $\det(I - VQK^{\text{ext}}V')_{L^2(\{t_1, \dots, t_n\} \times X)}$ and for $\det(I - QK^{\text{ext}})_{L^2(\{t_1, \dots, t_n\} \times X)}$. The same argument applies to the Fredholm determinant on the right-hand side of (20) by Assumption 3(iii), if we multiply it on the left by U_{t_1} and on the right by U'_{t_1} . Hence both sides of (20) are well-defined and one should really read the equality as

$$\begin{aligned} \det(I - VQK^{\text{ext}}V')_{L^2(\{t_1, \dots, t_n\} \times X)} \\ = \det(I - U_{t_1}(K_{t_1} - \overline{Q}_{t_1} \mathcal{W}_{t_1, t_2} \cdots \mathcal{W}_{t_{n-1}, t_n} \overline{Q}_{t_n} \mathcal{W}_{t_n, t_1} K_{t_1})U'_{t_1})_{L^2(X)}. \end{aligned}$$

Proof of Theorem 3.3. The proof of this result is a generalization of the proof of Theorem 1 of [40] (see also the Appendix of [37]). We will retain most of the notation of [37, 40], and as in those papers we use sans-serif fonts (e.g. W) for operators on $L^2(\{t_1, \dots, t_n\} \times X)$. This space can be identified with the space $\bigoplus_{t \in \{t_1, \dots, t_n\}} L^2(X)$, and hence we may (and will) think of an operator W on $L^2(\{t_1, \dots, t_n\} \times X)$ as an operator-valued $n \times n$ matrix. We will use serif fonts for the matrix entries (e.g. $W_{i,j} = W$ for some W acting on $L^2(X)$). All determinants throughout this proof are computed on $L^2(\{t_1, \dots, t_n\} \times X)$ unless otherwise indicated.

We will use repeatedly the following facts about trace class operators and Fredholm determinants on a separable Hilbert space \mathcal{H} :

- (i) If $A, B \in \mathcal{B}_1(\mathcal{H})$ then $AB \in \mathcal{B}_1(\mathcal{H})$ and

$$\det((I + A)(I + B))_{\mathcal{H}} = \det(I + A)_{\mathcal{H}} \det(I + B)_{\mathcal{H}}.$$

Moreover, if A and B are bounded linear operators on \mathcal{H} and both $AB, BA \in \mathcal{B}_1(\mathcal{H})$ then

$$\det(I + AB)_{\mathcal{H}} = \det(I + BA)_{\mathcal{H}}. \quad (21)$$

- (ii) An operator acting on $\bigoplus_{t \in \{t_1, \dots, t_n\}} \mathcal{H}$ is trace class if and only if all of its matrix entries are trace class.

To simplify notation throughout the proof we will replace subscripts of the form t_i by i , so for example $\mathcal{W}_{i,j} = \mathcal{W}_{t_i, t_j}$.

Recall that we are assuming $t_1 < t_2 < \dots < t_n$. Let $K = QK^{\text{ext}}$. Then K can be written as

$$K = Q(W^- K^d + W^+ (K^d - I)), \quad (22)$$

where

$$K_{ij}^d = K_i \mathbf{1}_{i=j}, \quad Q_{i,j} = Q_i \mathbf{1}_{i=j}$$

and W^- , W^+ are lower triangular, respectively strictly upper triangular, and defined by

$$W_{ij}^- = \mathcal{W}_{i,j} \mathbf{1}_{i \geq j}, \quad W_{ij}^+ = \mathcal{W}_{i,j} \mathbf{1}_{i < j}.$$

Here we are slightly abusing notation, because $\mathcal{W}_{i,j}$ is not defined for $i > j$. However, since W^- appears applied after K^d , the formula makes sense, with $[W^- K^d]_{i,j} = \mathcal{W}_{i,j} K_j$ for $i > j$. We also define the diagonal operators V , V' , U and U' by

$$V_{i,j} = V_i \mathbf{1}_{i=j}, \quad V'_{i,j} = V'_i \mathbf{1}_{i=j}, \quad U_{i,j} = U_i \mathbf{1}_{i=j} \quad \text{and} \quad U'_{i,j} = U'_i \mathbf{1}_{i=j}.$$

In order to manipulate the Fredholm determinant of $I - K$ we will need to make sure at each step that the appropriate operators preserve $L^2(X)$ and are trace class in $L^2(X)$ as needed. As a consequence, the proof is slightly cumbersome, so we will first briefly explain the main idea, ignoring some details and all analytical issues.

Our goal is to manipulate the determinant of $I - K$ in such a way that we end up with the determinant of an operator-valued matrix $I - \tilde{K}$ where only the first column of \tilde{K} is non-zero. If we achieve this, then we will have $\det(I - K) = \det(I - \tilde{K}) = \det(I - \tilde{K}_{1,1})_{L^2(X)}$, and all we will need to do is compute $K_{1,1}$. The key to obtain such an identity is the following observation. Using the semigroup property in Assumption 2(ii) one can check directly that

$$[(I + W^+)^{-1}]_{i,j} = I \mathbf{1}_{j=i} - \mathcal{W}_{i,i+1} \mathbf{1}_{j=i+1}. \quad (23)$$

This identity is meant in the sense of products of integral kernels, where the product of the identity operator with an integral kernel is defined in the obvious way. Now using the identity $\mathcal{W}_{i,j-1} K_{j-1} \mathcal{W}_{j-1,j} = \mathcal{W}_{i,j} K_j$ from Assumptions 2(ii) and 2(iii) we get that

$$[(W^- + W^+) K^d (I + W^+)^{-1}]_{i,j} = \mathcal{W}_{i,j} K_j - \mathcal{W}_{i,j-1} K_{j-1} \mathcal{W}_{j-1,j} \mathbf{1}_{j>1} = \mathcal{W}_{i,1} K_1 \mathbf{1}_{j=1}. \quad (24)$$

Note that only the first column of this matrix has non-zero entries. To take advantage of this fact we rewrite K as

$$K = Q(W^- + W^+) K^d (I + W^+)^{-1} (I + W^+) - QW^+, \quad (25)$$

so that

$$I - K = (I + QW^+) [I - (I + QW^+)^{-1} Q(W^- + W^+) K^d (I + W^+)^{-1} (I + W^+)].$$

The invertibility of $I + QW^+$ follows from the fact that QW^+ is strictly upper triangular. This fact also implies that $\det(I + QW^+) = 1$, and hence

$$\begin{aligned} \det(I - K) &= \det(I - (I + QW^+)^{-1} Q(W^- + W^+) K^d (I + W^+)^{-1} (I + W^+)) \\ &= \det(I - (I + W^+) (I + QW^+)^{-1} Q(W^- + W^+) K^d (I + W^+)^{-1}), \end{aligned}$$

where we have used the cyclic property of the determinant. Recalling that only the first column of $(W^- + W^+) K^d (I + W^+)^{-1}$ is non-zero we deduce that

$$\tilde{K} = (I + W^+) (I + QW^+)^{-1} Q(W^- + W^+) K^d (I + W^+)^{-1}$$

has the same property and hence $\det(I - K) = \det(I - \tilde{K}) = \det(I - \tilde{K}_{1,1})_{L^2(X)}$ as desired.

The rest of the proof will consist in making the above argument rigorous and precise and then computing the resulting $\tilde{K}_{1,1}$. Recall that, by Assumption 3(ii), each entry in the operator-valued matrix VKV' is trace class in $L^2(X)$. Let

$$W_1 = VQW^+V' \quad \text{and} \quad W_2 = VQ(W^- + W^+)K^dV'. \quad (26)$$

Since VQW^+V' is strictly upper triangular, we have $(VQW^+V')^{n+1} = 0$, so $I + W_1$ is invertible:

$$(I + W_1)^{-1} = \sum_{k=0}^n (-1)^k (VQW^+V')^k. \quad (27)$$

Therefore we can write

$$\det(I - VKV') = \det((I + W_1)(I - (I + W_1)^{-1}W_2)).$$

We remark that W_1 , W_2 and $(I + W_1)^{-1}$ are trace class in $L^2(X)$ by Assumption 3(ii) and (27), and thus from the last identity we deduce that

$$\det(I - VKV') = \det(I + W_1) \det(I - (I + W_1)^{-1}W_2) = \det(I - (I + W_1)^{-1}W_2), \quad (28)$$

where the second equality follows from the fact that, since W_1 is strictly upper triangular, its only eigenvalue is 0, so $\det(I + W_1) = 1$.

Write

$$(I + W_1)^{-1}W_2 = W_3W_4 \quad (29)$$

with

$$W_3 = (I + W_1)^{-1}VQ(W^- + W^+)K^d(I + W^+)^{-1}U' \quad \text{and} \quad W_4 = U(I + W^+)V'.$$

Here we are using (24) and the identity $KUU' = K$. We have already checked that W_3W_4 is trace class in $L^2(X)$. Thus if we prove that W_4W_3 is also trace class we can deduce from (21), (28) and (29) that

$$\det(I - VKV') = \det(I - W_4W_3). \quad (30)$$

We want to obtain an explicit expression for the kernel W_4W_3 . Note that, in view of (27) and the fact that $V'VQ = Q$, $V'(I + W_1)^{-1}VQ = (I + QW^+)^{-1}Q$, so all the factors V and V' cancel in W_4W_3 :

$$W_4W_3 = U(I + W^+)(I + QW^+)^{-1}Q(W^- + W^+)K^d(I + W^+)^{-1}U'.$$

From (24) and the semigroup property we deduce that, for $0 \leq k \leq n - i$,

$$\begin{aligned} & \left[(QW^+)^k Q(W^- + W^+)K^d(I + W^+)^{-1} \right]_{i,1} \\ &= \sum_{i < a_1 < \dots < a_k \leq n} Q_i \mathcal{W}_{i,a_1} Q_{a_1} \mathcal{W}_{a_1,a_2} \dots Q_{a_{k-1}} \mathcal{W}_{a_{k-1},a_k} Q_{a_k} \mathcal{W}_{a_k,1} K_1, \end{aligned}$$

while for $k > n - i$ the left-hand side above equals 0 (the case $k = 0$ is interpreted as $Q_i \mathcal{W}_{i,1} K_1$). Summing the above times $(-1)^k$ from $k = 0$ to $k = n - i$ we get directly from the last formula and (27) that

$$\begin{aligned} & \left[(I + QW^+)^{-1} Q(W^- + W^+)K^d(I + W^+)^{-1} \right]_{i,j} \\ &= \mathbf{1}_{j=1} \left[Q_i \mathcal{W}_{i,1} K_1 + \sum_{k=1}^{n-i} (-1)^k \sum_{i < a_1 < \dots < a_k \leq n} Q_i \mathcal{W}_{i,a_1} Q_{a_1} \mathcal{W}_{a_1,a_2} \dots Q_{a_{k-1}} \mathcal{W}_{a_{k-1},a_k} Q_{a_k} \mathcal{W}_{a_k,1} K_1 \right]. \end{aligned} \quad (31)$$

Note that only the first column of the above matrix contains non-zero entries. Since $U(I+W^+)$ is upper triangular and U' is diagonal, the same is true for W_4W_3 . Pre-multiplying (31) by $U(I+W^+)$ we get

$$(W_4W_3)_{i,1} = \sum_{j=i}^n \sum_{k=0}^{n-j} (-1)^k \sum_{j=a_0 < a_1 < \dots < a_k \leq n} U_i \mathcal{W}_{i,j} Q_j \mathcal{W}_{j,a_1} Q_{a_1} \mathcal{W}_{a_1,a_2} Q_{a_2} \mathcal{W}_{a_2,a_3} Q_{a_3} \mathcal{W}_{a_3,a_4} Q_{a_4} \mathcal{W}_{a_4,a_5} Q_{a_5} \mathcal{W}_{a_5,a_6} Q_{a_6} \mathcal{W}_{a_6,a_7} Q_{a_7} \mathcal{W}_{a_7,a_8} Q_{a_8} \mathcal{W}_{a_8,a_9} Q_{a_9} \mathcal{W}_{a_9,a_{10}} Q_{a_{10}} \mathcal{W}_{a_{10},1} K_1 U'_1. \quad (32)$$

By Lemma 3.1 we deduce that

$$(W_4W_3)_{i,1} = U_i [\mathcal{W}_{i,1} K_1 - \overline{Q}_i \mathcal{W}_{i,2} \overline{Q}_2 \dots \mathcal{W}_{n-1,n} \overline{Q}_n \mathcal{W}_{n,1} K_1] U'_1. \quad (33)$$

By Assumption 3(iii) this operator is trace class, which provides the needed justification for writing (30), and then since only the first column of W_4W_3 is non-zero we deduce that

$$\det(I - VKV') = \det(I - (W_4W_3)_{1,1})_{L^2(\mathbb{R})}.$$

Setting $i = 1$ in (33) yields the result. \square

In order to finish the proof of Theorem 3.3 it remains to prove Lemma 3.1.

Proof of Lemma 3.1. We start with the right-hand side of the identity. Replace each Q_i by $I - \overline{Q}_i$ except for the first one to get

$$\sum_{j=i}^n \sum_{k=0}^{n-j} (-1)^k \sum_{m=0}^k \binom{n-j-m}{k-m} (-1)^m \sum_{j=b_0 < b_1 < \dots < b_m \leq n} \mathcal{W}_{i,b_0} Q_{b_0} \mathcal{W}_{b_0,b_1} \dots \overline{Q}_{b_{m-1}} \mathcal{W}_{b_{m-1},b_m} \overline{Q}_{b_m} \mathcal{W}_{b_m,1} K_1$$

where, as in the above proof, we have written i instead of t_i in the subscripts. Interchanging the order of summation leads to $\sum_{j=i}^n \sum_{m=0}^{n-j} \sum_{k=m}^{n-j} (-1)^{k+m} \binom{n-j-m}{k-m} (\star)$, where (\star) represents the last sum above, and is independent of k . Noting that $\sum_{k=m}^{n-j} \binom{n-j-m}{k-m} (-1)^{k+m} = \mathbf{1}_{m=n-j}$, the above expression can be rewritten as

$$\begin{aligned} & \sum_{j=i}^n \sum_{j=b_0 < b_1 < \dots < b_{n-j} \leq n} \mathcal{W}_{i,b_0} Q_{b_0} \mathcal{W}_{b_0,b_1} \overline{Q}_{b_1} \dots \mathcal{W}_{b_{n-1},b_n} \overline{Q}_{b_n} \mathcal{W}_{b_n,1} K_1 \\ &= \sum_{j=i}^n \mathcal{W}_{i,j} (I - \overline{Q}_j) \mathcal{W}_{j,j+1} \overline{Q}_{j+1} \dots \mathcal{W}_{b_{n-1},b_n} \overline{Q}_{b_n} \mathcal{W}_{n,1} K_1 \\ &= \sum_{j=i}^n \left[\mathcal{W}_{i,j+1} \overline{Q}_{j+1} \mathcal{W}_{j+1,j+2} \overline{Q}_{j+2} \dots \mathcal{W}_{b_{n-1},b_n} \overline{Q}_{b_n} \mathcal{W}_{n,1} K_1 \right. \\ & \quad \left. - \mathcal{W}_{i,j} \overline{Q}_j \mathcal{W}_{j,j+1} \overline{Q}_{j+1} \dots \mathcal{W}_{b_{n-1},b_n} \overline{Q}_{b_n} \mathcal{W}_{n,1} K_1 \right] \\ &= \mathcal{W}_{i,1} K_1 - \overline{Q}_i \mathcal{W}_{i,2} \overline{Q}_2 \dots \mathcal{W}_{n-1,n} \overline{Q}_n \mathcal{W}_{n,1} K_1, \end{aligned}$$

where the last equality follows by telescoping. \square

4. A FEW EXAMPLES

We will now show how to apply Theorem 3.3 to a few examples of Fredholm determinants which arise in describing objects of interest in random matrix theory, growth processes, particle systems, tilings and representation theory. Our examples include extended determinantal point processes such as the stationary (GUE) Dyson Brownian motion, the Airy₂ process, and the Pearcey process; all of which are limits of ensembles of non-intersecting directed paths on weights graphs. We also include an extended determinant point process given by Markov processes on partitions related to the z -measures; this ensemble is *not* a

limit of a graph-based ensemble of non-intersecting directed paths. We also show how the identity applies to signed extended determinantal point processes such as the Airy_1 and $\text{Airy}_{2 \rightarrow 1}$ processes.

The proofs of the results in this section are postponed to the appendix.

4.1. Stationary (GUE) Dyson Brownian motion. Consider the eigenvalues of an $N \times N$ Hermitian matrix with each (algebraically independent) entry diffusing according to a stationary Ornstein-Uhlenbeck process (real valued on the diagonal and complex valued off the diagonal). The eigenvalues of this process are real valued and themselves form a Markov process, called the stationary Dyson Brownian motion. Its stationary marginal distribution is the $N \times N$ Gaussian Unitary Ensemble (GUE) eigenvalue distribution. We will consider this Dyson Brownian motion process in stationarity and write the i^{th} largest eigenvalue at time t as $\lambda_N(i; t)$. The collection of eigenvalues at time t is written $\lambda_N(\cdot; t) = (\lambda_N(1; t), \dots, \lambda_N(N; t))$ and the curve traced out by the i^{th} eigenvalue over time is written as $\lambda_N(i; \cdot)$. Then the graphs of $\lambda_N(1; \cdot), \dots, \lambda_N(N; \cdot)$ form an ensemble of non-intersecting curves (see, for example, Section 4.3.1 of [4]). This ensemble of curves is indexed by time t and curve label i and hence can be thought of as a random variable taking values in the space of continuous curves from $\{1, \dots, N\} \times \mathbb{R}$ to \mathbb{R} . We will write \mathbb{E} as the expectation operator for this random variable.

Definition 4.1. For times $t_1 < t_2 < \dots < t_n$ consider functions $q_{t_i} : \mathbb{R} \rightarrow \mathbb{R}$ and let $\bar{q}_{t_i}(x) = 1 - q_{t_i}(x)$. For a curve $g : \mathbb{R} \rightarrow \mathbb{R}$ define the functional \bar{q} by $\bar{q}(g) = \prod_{i=1}^n \bar{q}_{t_i}(g(t_i))$. One likewise defines the functional $q(g) = \prod_{i=1}^n q_{t_i}(g(t_i))$.

The stationary (GUE) Dyson Brownian motion is an extended determinantal point process. In particular this means that for any functions q_{t_i} (as above),

$$\mathbb{E} \left[\prod_{j=1}^N \bar{q}(\lambda_N(j; \cdot)) \right] = \det(I - Q K_{\text{GUE}, N}^{\text{ext}})_{L^2(\{t_1, \dots, t_n\} \times \mathbb{R})} \quad (34)$$

as long as both sides are well-defined, where Q is defined as in (19) and $K_{\text{GUE}, N}^{\text{ext}}$ is the *extended Hermite kernel* (see e.g. [45]):

$$K_{\text{GUE}, N}^{\text{ext}}(s, x; t, y) = \begin{cases} \sum_{k=0}^{N-1} e^{k(s-t)} \varphi_k(x) \varphi_k(y) & \text{if } s \geq t, \\ - \sum_{k=N}^{\infty} e^{k(s-t)} \varphi_k(x) \varphi_k(y) & \text{if } s < t. \end{cases}$$

Here $\varphi_k(x) = e^{-x^2/2} p_k(x)$ and p_k is the k -th normalized Hermite polynomial (so that $\|\varphi_k\|_2 = 1$).

Writing

$$D = -\frac{1}{2}(\Delta - x^2 + 1),$$

the *harmonic oscillator functions* φ_k satisfy $D\varphi_k = k\varphi_k$. Then the *Hermite kernel*

$$K_{\text{GUE}, N}(x, y) = K_{\text{GUE}, N}(0, x; 0, y) = \sum_{k=0}^{N-1} \varphi_k(x) \varphi_k(y)$$

acts as the projection operator onto $\text{span}\{\varphi_0, \dots, \varphi_{N-1}\}$. In the notation of Theorem 3.3 we are taking $X = \mathbb{R}$, μ the Lebesgue measure, and for $1 \leq i < j \leq n$

$$\mathcal{W}_{t_i, t_j}(x, y) = e^{-(t_j - t_i)D}(x, y) = \sum_{k=0}^{\infty} e^{-(t_j - t_i)k} \varphi_k(x) \varphi_k(y), \quad K_{t_i} = K_{\text{GUE}, N},$$

$$\mathcal{W}_{t_j, t_i} K_{t_i}(x, y) = e^{(t_j - t_i)D} K_{\text{GUE}, N}(x, y) = \sum_{k=0}^{N-1} e^{(t_j - t_i)k} \varphi_k(x) \varphi_k(y).$$

Applying Theorem 3.3 we conclude:

Corollary 4.2. *Fix $t_1 < \dots < t_n$ and write $\tau = \min_{i=1, \dots, n-1} |t_{i+1} - t_i|$. For each $1 \leq i \leq n$ choose a function $q_{t_i} \in L^1_{\text{loc}}(\mathbb{R})$ satisfying $\sup_{x \in \mathbb{R}} e^{-\kappa x^2} |q_{t_i}(x)| < \infty$ for some $\kappa \in (0, \frac{1}{2\sqrt{2}} \tanh(\tau/\sqrt{2}))$. Then*

$$\mathbb{E} \left[\prod_{j=1}^N q(\lambda_j^N) \right] = \det(I - K_{\text{GUE}, N} + Q_{t_1} e^{(t_1 - t_2)D} Q_{t_2} \dots Q_{t_n} e^{(t_n - t_1)D} K_{\text{GUE}, N})_{L^2(\mathbb{R})}.$$

Note that we have removed the bars over q and Q by replacing q_{t_i} 's by $(1 - q_{t_i})$'s.

4.1.1. Continuum statistics. We may now take a continuous time limit of the above formula (in the style of [22]). Consider a function $h : \mathbb{R} \times \mathbb{R} \rightarrow [0, \infty]$ and $\ell < r$. Define an operator $\Gamma_{\ell, r}^h$ acting on $L^2(\mathbb{R})$ as follows: $\Gamma_{\ell, r}^h f(\cdot) = u(r, \cdot)$, where $u(r, \cdot)$ is the solution at time r of

$$\partial_t u = -Du - hu \quad (35)$$

with initial data $u(\ell, x) = f(x)$. By the Feynman-Kac formula we may also express the action of this operator in terms of a path-integral through a potential h as

$$\Gamma_{\ell, r}^h f(x) = \mathbb{E}_{b(\ell)=x} \left[f(b(r)) e^{-\frac{1}{2} \int_{\ell}^r (2h(s, b(s)) + b(s)^2 - 1) ds} \right] \quad (36)$$

where the expectation is over a (standard) Brownian motion $b(\cdot)$ started at time ℓ with $b(\ell) = x$ and run until time r .

Let $t_1 = \ell$, $t_n = r$ and the t_i be spaced equally in between with step size $\delta = (r - \ell)/(n - 1)$. Then letting $q_{t_i}(x) = 1 - \delta h(t_i, x)$ and taking $n \rightarrow \infty$ the above formula yields:

Proposition 4.3. *For any interval $[\ell, r]$ and continuous bounded function $h : \mathbb{R} \times \mathbb{R} \rightarrow [0, \infty]$*

$$\mathbb{E} \left[\prod_{j=1}^N \exp \left(- \int_{\ell}^r h(t, \lambda_N(j; t)) dt \right) \right] = \det(I - K_{\text{GUE}, N} + \Gamma_{\ell, r}^h e^{(t_n - t_1)D} K_{\text{GUE}, N})_{L^2(\mathbb{R})}. \quad (37)$$

Remark 4.4. The condition on h is not optimal, but it makes the arguments simpler. A different class of functions h for which the result holds is the following. Fix a function $g \in H^1([\ell, r])$ and set $h(t, x) = 0$ for $x < g(t)$ and infinity otherwise. Then the left-hand side of (37) becomes $\mathbb{P} \left[\bigcap_{j=1}^N \{ \lambda_N(j; t) < g(t) \forall t \in [\ell, r] \} \right]$ and the right-hand side makes perfect sense as well, with $\Gamma_{\ell, r}^h$ now being the solution operator of a certain boundary operator involving g . This case corresponds to calculating the probability that on the entire interval $[\ell, r]$, the top curve of the Dyson Brownian motion remains below the function $g(t)$. This is the same type of result shown in [22] for the Airy₂ process, and the proof for this case can be easily adapted from the arguments in that paper.

4.1.2. *Rescaled process.* Now introduce the rescaled process

$$\tilde{\lambda}_N(i; t) = \sqrt{2}N^{1/6}(\lambda_N(i; N^{-1/3}t) - \sqrt{2N}).$$

Changing variables $x \mapsto \frac{1}{\sqrt{2}N^{1/6}}x + \sqrt{2N}$, $y \mapsto \frac{1}{\sqrt{2}N^{1/6}}y + \sqrt{2N}$ in the kernel accordingly, we immediately obtain:

Corollary 4.5. *For any $t_1 < \dots < t_n$ and functions $q_{t_i} : \mathbb{R} \rightarrow \mathbb{R}$, $1 \leq i \leq n$, satisfying the same conditions as in Corollary 4.2, we have*

$$\mathbb{E} \left[\prod_{j=1}^N q(\tilde{\lambda}_N(j; \cdot)) \right] = \det(I - \tilde{K}_{\text{GUE}, N} + Q_{t_1} e^{(t_1 - t_2)H_N} Q_{t_2} \dots Q_{t_n} e^{(t_n - t_1)H_N} \tilde{K}_{\text{GUE}, N})_{L^2(\mathbb{R})}. \quad (38)$$

where the kernel of $\tilde{K}_{\text{GUE}, N}$ is given by

$$\tilde{K}_{\text{GUE}, N}(x, y) = \frac{1}{\sqrt{2}N^{1/6}} K_{\text{GUE}, N} \left(\frac{x}{\sqrt{2}N^{1/6}} + \sqrt{2N}, \frac{y}{\sqrt{2}N^{1/6}} + \sqrt{2N} \right)$$

and the operator

$$H_N = -\Delta + x + \frac{x^2}{2N^{2/3}}.$$

The above rescaling corresponds to focusing in on the top curves of the Dyson Brownian motion. In the limit N goes to infinity, $\tilde{K}_{\text{GUE}, N}$ converges to the Airy_2 kernel K_2 and H_N converges to the Airy Hamiltonian H (defined in the Introduction and below in Section 4.2). So in the limit as N goes to infinity we recover the formula for the Airy_2 process as expected. The operator in the Fredholm determinant in the right-hand side of (38) converges in trace class to the corresponding one with K_2 and H , which means that all of the left-hand side probabilities have limits. This can certainly be proved under some additional (though not optimal) assumptions on the q_{t_i} as in Corollary 4.6, but we choose to treat the Airy_2 process independently.

4.2. The Airy_2 line ensemble. The multi-layer Airy_2 process [26, 36] is the limit of the stationary (GUE) Dyson Brownian motion under the scaling of Section 4.1.2. In particular for $t \in \mathbb{R}$ consider the point process corresponding to $\{\tilde{\lambda}_N(i; t) : 1 \leq i \leq N\}$. As N goes to infinity, this point process converges in the vague topology to a limiting point process with an infinite number of simple points which we write as $\{\text{Airy}_2(i; t) : i \in \mathbb{Z}_{\geq 1}\}$ (labeled so that $\text{Airy}_2(i; t) > \text{Airy}_2(j; t)$ for $i < j$). This convergence can be strengthened so that for any fixed set $t_1 < t_2 < \dots < t_n$, the n -tuple of $\tilde{\lambda}$ -point processes has a limit $\{\text{Airy}_2(i; t) : i \in \mathbb{Z}_{\geq 1}, t \in \{t_1, \dots, t_n\}\}$. This limiting collection of point processes is consistent and can be completed to a point process valued stochastic process indexed by $t \in \mathbb{R}$. This process is called the multi-layer Airy_2 process. As it is the limit of a stationary (in t) process, it is also stationary.

There exists a continuous version of this process [21] so that Airy_2 can be thought of as a random variable taking values in the space of $\mathbb{Z}_{\geq 1}$ indexed, continuous and non-intersecting curves from \mathbb{R} to \mathbb{R} . The convention is that $\text{Airy}_2(1; \cdot)$ represents the top curve (i.e., the limit of $\tilde{\lambda}_N(1; \cdot)$). The continuous version of the multi-layer Airy_2 process is called the Airy_2 line ensemble.

Since the Dyson Brownian motion was an extended determinantal point process (34), so too is the multi-layer Airy_2 process. Analogous to (34), and with the functional \bar{q} given in

Definition 4.1 and operator Q given in (19),

$$\mathbb{E} \left[\prod_{j=1}^{\infty} \bar{q}(\text{Airy}_2(j; \cdot)) \right] = \det(I - QK_2^{\text{ext}})_{L^2(\{t_1, \dots, t_n\} \times \mathbb{R})} \quad (39)$$

where K_2^{ext} is the *extended Airy₂ kernel*

$$K_2^{\text{ext}}(s, x; t, y) = \begin{cases} \int_0^{\infty} d\lambda e^{-\lambda(s-t)} \text{Ai}(x + \lambda) \text{Ai}(y + \lambda) & \text{if } s \geq t, \\ - \int_{-\infty}^0 d\lambda e^{-\lambda(s-t)} \text{Ai}(x + \lambda) \text{Ai}(y + \lambda) & \text{if } s < t, \end{cases} \quad (40)$$

and $\text{Ai}(\cdot)$ is the Airy function. In order for the above expectation to make sense, one has to impose conditions on the functions q_{t_i} , such as in Corollary 4.6.

To put this example in the setting of Theorem 3.3 we take $X = \mathbb{R}$, μ the Lebesgue measure, and consider the *Airy Hamiltonian* defined as

$$H = -\Delta + x.$$

H has the shifted Airy functions $\text{Ai}_{\lambda}(x) = \text{Ai}(x - \lambda)$ as its generalized eigenfunctions: $H\text{Ai}_{\lambda}(x) = \lambda\text{Ai}_{\lambda}(x)$. Define the Airy_2 kernel K_2 as the projection of H onto its negative generalized eigenspace:

$$K_2(x, y) = \int_0^{\infty} d\lambda \text{Ai}(x + \lambda) \text{Ai}(y + \lambda).$$

Then it is not hard to check that, in the notation of Theorem 3.3, (40) corresponds to taking, for $1 \leq i < j \leq n$,

$$\begin{aligned} \mathcal{W}_{t_i, t_j}(x, y) &= e^{-(t_j - t_i)H}(x, y) = \int_{-\infty}^{\infty} d\lambda e^{\lambda(t_j - t_i)} \text{Ai}(x + \lambda) \text{Ai}(y + \lambda), \quad K_{t_i} = K_2, \\ \mathcal{W}_{t_j, t_i} K_{t_i}(x, y) &= e^{(t_j - t_i)H} K_2(x, y) = \int_0^{\infty} d\lambda e^{-\lambda(t_j - t_i)} \text{Ai}(x + \lambda) \text{Ai}(y + \lambda). \end{aligned}$$

Note that \mathcal{W}_{t_i, t_j} is only well-defined on the range of K_2 . Applying Theorem 3.3 allows to conclude:

Corollary 4.6. *Fix $t_1 < \dots < t_n$ and let $\tau = \min_{i=1, \dots, n-1} \{t_{i+1} - t_i\}$. Choose functions $q_{t_i} \in L^1_{\text{loc}}(\mathbb{R})$, $1 \leq i \leq n$, such that $\sup_{x \geq 0} e^{-rx} |q_{t_i}(x)| < \infty$ for some $0 < r < \tau$ and $\sup_{x < 0} \varphi(x) |1 - q_{t_i}(x)| < \infty$ for some function $\varphi(x)$ such that $\int_{-\infty}^0 dx e^{-2(t_n - t_1)x} \varphi(x)^{-2} < \infty$, $1 \leq i \leq n$. Then*

$$\mathbb{E} \left[\prod_{j=1}^{\infty} q(\text{Airy}_2(j; \cdot)) \right] = \det(I - K_2 + Q_{t_1} e^{(t_1 - t_2)H} Q_{t_2} \dots Q_{t_n} e^{(t_n - t_1)H} K_2)_{L^2(\mathbb{R})}.$$

This formula is also the limit of the right-hand side of (38) as N goes to infinity.

Since the Airy line ensemble is a continuous version of the multi-layer Airy_2 process, we may take a continuum limit of the above formula, in the same manner as done in Section 4.1.1. The PDE which $\Gamma_{\ell, r}^h$ is solving is now $\partial_t u = -Hu - hu$ (corresponding to replacing D by H in (35)) and the result is that for any interval $[\ell, r]$ and suitable function $h : \mathbb{R} \times \mathbb{R} \rightarrow [0, \infty]$ (for example h can be taken to be bounded, continuous, and such that $h(t, x) = 0$ for any $t \in [\ell, r]$ and $x < M$ for some $M \in \mathbb{R}$),

$$\mathbb{E} \left[\prod_{j=1}^{\infty} \exp \left(- \int_{\ell}^r h(t, \text{Airy}_2(j; t)) dt \right) \right] = \det(I - K_2 + \Gamma_{\ell, r}^h e^{(r - \ell)H} K_2)_{L^2(\mathbb{R})}. \quad (41)$$

We will omit the proof of this statement, which can be adapted from the proofs of Proposition 4.3 and Corollary 4.6 together with the proof of Proposition 3.2 in [22]. Taking $h(t, x)$ to be 0 for $x < g(t)$ and infinity otherwise we recover Theorem 2 of [22].

4.3. The Pearcey process. There are many other multi-layer processes which arise as scaling limits of non-intersecting ensembles of Brownian motions (or similar diffusions) for which we can apply Theorem 3.3 (see for instance Airy-like processes [2, 3, 6, 16]; bulk limits such as the Sine process [45], Pearcey process [46] or Tacnode process [8]; hard edge limits like the Bessel process [30]).

To illustrate this point we will show how a Fredholm determinant involving the Pearcey kernel can be rewritten via Theorem 3.3.

Let us briefly and informally recall one way the Pearcey process arises as a scaling limit of Brownian bridges. Consider $2N$ Brownian bridges on the time interval $[-N, N]$ such that all $2N$ of them start at height 0 and N of them end at height b and the other N end at height $-b$. Condition these Brownian bridges not to intersect (as can be done by spacing their starting and ending points by ε and letting ε go to zero). When $b = 0$ the limit shape of the ensemble of conditioned Brownian bridges has a limit shape which is elliptical (and the ensemble is sometimes called a watermelon) and the fluctuations around the top of this limit shape are described (in the limit as N goes to infinity) by the Airy_2 line ensemble minus a parabolic shift.

When the endpoints parameter $b = cN$, the limit shape has a cusp at some time $t = c'N$, where $c' \in (-1, 1)$ is a function of c . For $t_1 < t_2 < \dots < t_n$, the N -tuple of point processes formed by the heights (properly centered and normalized by $N^{1/4}$ near the height of the cusp) of the Brownian bridges at times $c'N + t_i N^{1/2}$, $1 \leq i \leq n$, converges in the vague topology as N goes to infinity to a limit which is called the *Pearcey process*, \mathcal{P} , see [5, 19, 20, 34, 46]. It is a point process valued stochastic process indexed by $t \in \mathbb{R}$. At each time t the point process can be indexed by \mathbb{Z} as $\{\mathcal{P}(j; t) : j \in \mathbb{Z}\}$.

Analogously to (34), and with the functional \bar{q} given in Definition 4.1 and operator Q given in (19),

$$\mathbb{E} \left[\prod_{j=-\infty}^{\infty} \bar{q}(\mathcal{P}(j; \cdot)) \right] = \det(I - Q K_{\text{PrC}}^{\text{ext}})_{L^2(\{t_1, \dots, t_n\} \times \mathbb{R})}, \quad (42)$$

where $K_{\text{PrC}}^{\text{ext}}$ is the *extended Pearcey kernel*

$$K_{\text{PrC}}^{\text{ext}}(s, x; t, y) = -\frac{1}{\sqrt{4\pi(t-s)}} \exp\left(-\frac{(y-x)^2}{4(t-s)}\right) \mathbf{1}_{t>s} \\ + \frac{1}{(2\pi i)^2} \int_C du \int_{-\infty}^{\infty} dv \frac{e^{-v^4/4+tv^2/2-yv}}{e^{-u^4/4+su^2/2-xu}} \frac{1}{v-u}, \quad (43)$$

and where C is the contour consisting of the rays going from $\pm\infty e^{i\pi/4}$ to 0 and from 0 to $\pm\infty e^{-i\pi/4}$.

In the setting of Theorem 3.3 we take $X = \mathbb{R}$, μ the Lebesgue measure, and for $t_i < t_j$ define

$$\mathcal{W}_{t_i, t_j} = e^{\frac{1}{2}(t_j - t_i)\Delta}, \quad K_{t_i}(x, y) = K_{\text{PrC}}^{t_i}(x, y) := K_{\text{PrC}}^{\text{ext}}(t_i, x; t_i, y), \\ \mathcal{W}_{t_j, t_i} K_{t_i}(x, y) = K_{\text{PrC}}^{\text{ext}}(t_j, x; t_i, y).$$

The semigroup property is obviously satisfied, while for $i < j$

$$\begin{aligned} \mathcal{W}_{t_i, t_j} K_{\text{Prc}}^{t_j}(x, y) &= \int_{-\infty}^{\infty} dz \frac{1}{\sqrt{2\pi(t_j - t_i)}} e^{-\frac{(x-z)^2}{2(t_j - t_i)}} \frac{1}{(2\pi i)^2} \int_C du \int_{-\infty}^{\infty} dv \frac{e^{-\frac{v^4}{4} + t_j \frac{v^2}{2} - yv}}{e^{-\frac{u^4}{4} + t_j \frac{u^2}{2} - zu}} \frac{1}{v - u} \\ &= \frac{1}{(2\pi i)^2} \int_C du \int_{-\infty}^{\infty} dv \frac{e^{-\frac{v^4}{4} + t_j \frac{v^2}{2} - yv}}{e^{-\frac{u^4}{4} + t_j \frac{u^2}{2} - xu}} \frac{1}{v - u} = K_{\text{Prc}}^{\text{ext}}(t_i, x; t_j, y) + \mathcal{W}_{t_i, t_j} = K_{\text{Prc}}^{t_i} \mathcal{W}_{t_i, t_j}(x, y), \end{aligned}$$

where the second equality follows from computing a simple Gaussian integral and the last equality is obtained similarly. Likewise one can check that for $i < j$ we have $\mathcal{W}_{t_i, t_j} \mathcal{W}_{t_j, t_i} K_{\text{Prc}}^{t_i} = K_{\text{Prc}}^{t_i}$. Hence Assumption 2 is satisfied, and from Theorem 3.3 we deduce the following:

Corollary 4.7. *For any $t_1 < t_2 < \dots < t_n$ and functions $q_{t_i} : \mathbb{R} \rightarrow \mathbb{R}$, $1 \leq i \leq n$ so that Assumptions 1 and 3 are satisfied, we have*

$$\mathbb{E} \left[\prod_{j=-\infty}^{\infty} q(\mathcal{P}(j; \cdot)) \right] = \det(I - K_{\text{Prc}}^{t_1} + Q_{t_1} e^{\frac{1}{2}(t_2 - t_1)\Delta} Q_{t_2} \dots e^{\frac{1}{2}(t_n - t_{n-1})\Delta} Q_{t_n} e^{\frac{1}{2}(t_1 - t_n)\Delta} K_{\text{Prc}}^{t_1})_{L^2(\mathbb{R})}. \quad (44)$$

In particular, the formula holds for the case $q_{t_i}(x) = \mathbf{1}_{x \leq a_i}$.

We do not attempt here to provide more general conditions on the functions q_{t_i} so that the formula holds.

4.4. The $\text{Airy}_{2 \rightarrow 1}$ and $\text{Airy}_{2 \rightarrow 1}$ processes. All of the examples considered thus far have involved probability measures on ensembles of non-intersecting paths or their scaling limits. Going back to the discrete setting of Theorem 2.2, there was no condition that the measure on non-intersecting paths be positive. This condition is not met, for example, in the case of the Airy_1 and $\text{Airy}_{2 \rightarrow 1}$ processes. These are real valued stochastic processes which are the scaling limits of marginals of measures (not entirely positive) on non-intersecting paths [9–11, 42]. Even though the ensemble measure is not entirely positive, the marginal is a probability measure.

We will focus on the $\text{Airy}_{2 \rightarrow 1}$ process obtained in [11], since a similar result to that which we now state has already shown up in [40]. The $\text{Airy}_{2 \rightarrow 1}$ process is a continuous time (non-stationary) real valued process $\text{Airy}_{2 \rightarrow 1} : \mathbb{R} \rightarrow \mathbb{R}$ given by its finite-dimensional distributions

$$\mathbb{P} \left(\bigcap_{k=1}^n \{ \text{Airy}_{2 \rightarrow 1}(t_k) \leq x_k \} \right) = \det(I - \chi K_{2 \rightarrow 1})_{L^2(\{t_1, \dots, t_n\} \times \mathbb{R})} \quad (45)$$

for $t_1 < \dots < t_n$, where $\chi f(t_i, x) = \mathbf{1}_{x \geq x_i} f(x)$ and

$$\begin{aligned} K_{2 \rightarrow 1}^{\text{ext}}(s, x; t, y) &= -\frac{1}{\sqrt{4\pi(t-s)}} \exp \left(-\frac{(\tilde{y} - \tilde{x})^2}{4(t-s)} \right) \mathbf{1}_{t > s} \\ &\quad + \frac{1}{(2\pi i)^2} \int_{\gamma_+} dw \int_{\gamma_-} dz \frac{e^{w^3/3 + tw^2 - \tilde{y}w}}{e^{z^3/3 + sz^2 - \tilde{x}z}} \frac{2w}{(z-w)(z+w)} \end{aligned} \quad (46)$$

with

$$\tilde{x} = x - (s^-)^2, \quad \tilde{y} = y - (t^-)^2,$$

notation $r^- = \min\{0, r\}$, and the paths γ_+, γ_- satisfying $-\gamma_+ \subseteq \gamma_-$ with $\gamma_+ : e^{i\phi_+}\infty \rightarrow e^{-i\phi_+}\infty$, $\gamma_- : e^{-i\phi_-}\infty \rightarrow e^{i\phi_-}\infty$ for some $\phi_+ \in (\pi/3, \pi/2)$, $\phi_- \in (\pi/2, \pi - \phi_+)$. The $\text{Airy}_{2 \rightarrow 1}$ process crosses over between the Airy_2 and the Airy_1 processes in the sense that $\text{Airy}_{2 \rightarrow 1}(t + \tau)$ converges to $2^{1/3} \text{Airy}_1(2^{-2/3}t)$ as $\tau \rightarrow \infty$ and to $\text{Airy}_2(1; t)$ (the

Airy₂ process, i.e. the top line of the multi-layer Airy₂ process) when $\tau \rightarrow -\infty$ (in the sense of finite dimensional distributions). It is expected to govern the asymptotic spatial fluctuations in random growth models when the initial conditions are deterministic near the point where the hydrodynamic profile changes from flat to curved. In particular, it is shown in [11] that it governs the asymptotic fluctuations near the profile switch point for the totally asymmetric simple exclusion process starting with particles only at the even negative integers.

We take again $X = \mathbb{R}$ and μ the Lebesgue measure, and for $i < j$ we define

$$\mathcal{W}_{t_i, t_j}(x, y) = e^{(t_j - t_i)\Delta}(x - (t_i^-)^2, y - (t_j^-)^2), \quad K_{t_i}(x, y) = K_{2 \rightarrow 1}^{t_i}(x, y) := K_{2 \rightarrow 1}^{\text{ext}}(t_i, x; t_i, y) \\ \mathcal{W}_{t_j, t_i} K_{t_i}(x, y) = K_{2 \rightarrow 1}^{\text{ext}}(t_j, x; t_i, y).$$

Proceeding as in Section 4.3 one checks that these choices satisfy Assumption 2, and hence (under the additional assumptions) we may apply Theorem 3.3. Using the translation invariance of the heat kernel to rearrange the shifts appearing in the resulting formula we get:

Corollary 4.8. *For any $t_1 < t_2 < \dots < t_n$, we have*

$$\mathbb{P}\left(\bigcap_{k=1}^n \{\text{Airy}_{2 \rightarrow 1}(t_k) \leq x_k\}\right) \\ = \det(I - K_{2 \rightarrow 1}^{t_1} + \bar{P}_{\tilde{x}_1} e^{(t_2 - t_1)\Delta} \bar{P}_{\tilde{x}_2} \dots e^{(t_n - t_{n-1})\Delta} \bar{P}_{\tilde{x}_n} e^{(t_1 - t_n)\Delta} K_{2 \rightarrow 1}^{t_1})_{L^2(\mathbb{R})}, \quad (47)$$

where $\tilde{x}_i = x_i - (t_i^-)^2$, $\bar{P}_a f(x) = \mathbf{1}_{x \leq a} f(x)$ and $K_{2 \rightarrow 1}^{t_1}(x, y) = K_{2 \rightarrow 1}^{\text{ext}}(t_1, x + (t_1^-)^2; t_1, y + (t_1^-)^2)$.

In the formula, $e^{(t_1 - t_n)\Delta} K_{2 \rightarrow 1}^{t_1}$ should be interpreted as $K_{2 \rightarrow 1}^{\text{ext}}(t_n, x + (t_n^-)^2; t_1, y + (t_1^-)^2)$. One can use this formula directly to recover the analogous path-integral kernel formulas for the Airy₁ and Airy₂ processes in the appropriate limits, and thus show that Airy_{2 \rightarrow 1} interpolates between these two processes.

4.5. Markov processes on partitions and z -measures. The z -measures are a remarkable family of probability distributions on partitions that arise in representation theory of the infinite-symmetric group. They can be viewed as determinantal point processes on the one-dimensional lattice with infinite many particles, and they degenerate to a variety of well-known discrete and continuous determinantal point processes, see [12, 13, 35] and references therein.

In [15], a Markov process on partitions that preserves the z -measures was constructed. Its dynamical correlation functions are determinantal, and they can be described via the corresponding extended kernel, see Section 6 of [15]. One particular limit of this Markov process can be seen as ‘space-like’ space-time sections of the multilayer polynuclear growth process of [36], see [14].

Note that it is not known how to obtain the z -measures and the corresponding Markov processes as a limit of an ensemble of nonintersecting paths. However, these objects can be viewed as an *analytic continuation* of an ensemble of nonintersecting birth-and-death processes *in the number of paths*, see Section 6.5 of [15].

By encoding a partition $\lambda = (\lambda_1 \geq \lambda_2 \geq \dots)$ by the point configuration $\{\lambda_i - i + \frac{1}{2}\}_{i \geq 1}$, the Markov process can be written as $\{\mathcal{Z}(j; t) : j \in \mathbb{Z}_{\geq 1}, t \in \mathbb{R}\}$. Here $\mathcal{Z}(j; t)$ takes values in $\mathbb{Z}' = \mathbb{Z} + \frac{1}{2}$.

Similarly to (34), and with the functional \bar{q} given in Definition 4.1 and operator Q given in (19),

$$\mathbb{E} \left[\prod_{j=1}^{\infty} \bar{q}(\mathcal{Z}(j; \cdot)) \right] = \det(I - QK_{z,z',\xi}^{\text{ext}})_{L^2(\{t_1, \dots, t_n\} \times \mathbb{Z}')} \quad (48)$$

where $K_{z,z',\xi}^{\text{ext}}$ is the *extended hypergeometric kernel* which we will now define.

For parameters $z, z' \in \mathbb{C}$ such that either $z' = \bar{z} \in \mathbb{C} \setminus \mathbb{Z}$ or $m < z, z' < m+1$ for a $m \in \mathbb{Z}$, and $\xi \in (0, 1)$ define a second order difference operator $D_{z,z',\xi}$ on \mathbb{Z}' , depending on (z, z', ξ) and acting on functions $f(\cdot) \in \ell^2(\mathbb{Z}')$ as follows

$$(D_{z,z',\xi}f)(x) = \sqrt{\xi(z+x+\frac{1}{2})(z'+x+\frac{1}{2})}f(x+1) + \sqrt{\xi(z+x-\frac{1}{2})(z'+x-\frac{1}{2})}f(x-1) - (x+\xi(z+z'+x))f(x). \quad (49)$$

This is a self-adjoint operator with discrete simple spectrum $(1-\xi)\mathbb{Z}'$. Its eigenfunctions ψ_a ,

$$D_{z,z',\xi}\psi_a = (1-\xi)a\psi_a,$$

are explicitly written through the Gauss hypergeometric function (see [15], equation (5.1)). We normalize them by the condition $\|\psi_a\|_{\ell^2(\mathbb{Z}')} = 1$. Then

$$K_{z,z',\xi}^{\text{ext}}(s, x; t, y) = \begin{cases} \sum_{a \in \mathbb{Z}'_+} e^{-a(s-t)} \psi_a(x) \psi_a(y) & \text{if } s \geq t, \\ - \sum_{a \in \mathbb{Z}'_-} e^{-a(s-t)} \psi_a(x) \psi_a(y) & \text{if } s < t, \end{cases} \quad (50)$$

where $\mathbb{Z}'_{\pm} = \{\pm\frac{1}{2}, \pm\frac{3}{2}, \pm\frac{5}{2}, \dots\}$.

Let $D'_{z,z',\xi} = -(1-\xi)^{-1}D_{z,z',\xi}$. Then in the setting of Theorem 3.3, (50) corresponds to taking $X = \mathbb{Z}'$, μ the counting measure and, for $i < j$,

$$\mathcal{W}_{t_i, t_j} = e^{(t_j - t_i)D'_{z,z',\xi}}, \quad K_{t_i}(x, y) = K_{z,z',\xi}(x, y) := K_{z,z',\xi}^{\text{ext}}(0, x; 0, y) = \sum_{a \in \mathbb{Z}'_+} \psi_a(x) \psi_a(y),$$

$$\mathcal{W}_{t_j, t_i} K_{t_i}(x, y) = K_{z,z',\xi}^{\text{ext}}(t_j, x; t_i, y) = \sum_{a \in \mathbb{Z}'_+} e^{-a(t_j - t_i)} \psi_a(x) \psi_a(y).$$

Thus from Theorem 3.3 we deduce the following:

Corollary 4.9. *For any $t_1 < t_2 < \dots < t_n$ and functions $q_{t_i} : \mathbb{Z}' \rightarrow \mathbb{R}$, $1 \leq i \leq n$, so that Assumptions 1 and 3 are satisfied, we have*

$$\mathbb{E} \left[\prod_{j=1}^{\infty} q(\mathcal{Z}(j; \cdot)) \right] = \det(I - K_{z,z',\xi} + Q_{t_1} e^{(t_2 - t_1)D'_{z,z',\xi}} Q_{t_2} \dots e^{(t_n - t_{n-1})D'_{z,z',\xi}} Q_{t_n} e^{(t_1 - t_n)D'_{z,z',\xi}} K_{z,z',\xi})_{L^2(\mathbb{Z}')} \quad (51)$$

Let us remark that if all the functions q_{t_i} have finite support then all the needed analytic assumptions are automatically satisfied because we are working in an L^2 space on a finite set. Of course, such a restriction is unnecessarily harsh, but we will not pursue this issue here any further.

APPENDIX A. PROOFS OF THE RESULTS FROM SECTION 4

We recall the following facts about trace class and Hilbert-Schmidt norms (see e.g. [43]) of operators in $L^2(X)$ for some measurable space (X, Σ, μ) , which we will use repeatedly without reference:

$$\|AB\|_1 \leq \|A\|_2 \|B\|_2, \quad \|AB\|_1 \leq \|A\|_{\text{op}} \|B\|_1,$$

and if A has integral kernel $A(x, y)$,

$$\|A\|_2 = \left(\int \mu(dx) \mu(dy) |A(x, y)|^2 \right)^{1/2},$$

for each A, B in the appropriate space, where $\|\cdot\|_{\text{op}}$ denotes the operator norm in $L^2(X)$.

Throughout this section c and c' will denote positive constants whose value may change from line to line.

Proof of Corollary 4.2. Checking Assumption 2 is straightforward. We will take in this case $V_{t_i} = V'_{t_i} = U_{t_i} = U'_{t_i} = I$, and thus Assumption 1 is contained in Assumption 3, which we check next.

Condition (i) is trivial. Given functions ψ_1 and ψ_2 write $\psi_1 \otimes \psi_2$ for the kernel $\psi_1(x)\psi_2(y)$ and let ϕ be any function with $\int \phi^2 = 1$. Then we can write $\bar{Q}_{t_i}(\varphi_k \otimes \varphi_k) = (\bar{Q}_{t_i} \varphi_k \otimes \phi)(\phi \otimes \varphi_k)$, so that $\|\bar{Q}_{t_i}(\varphi_k \otimes \varphi_k)\|_1 \leq \|\bar{Q}_{t_i} \varphi_k \otimes \phi\|_2 \|\phi \otimes \varphi_k\|_2$ (note that we need to consider the operators with the bars because of the remark following the statement of the corollary). Now slightly abusing notation to write $\|\cdot\|_2$ both for the Hilbert-Schmidt norm of operators in $L^2(\mathbb{R})$ and for the norm of this last space, we have

$$\|\bar{Q}_{t_i} \varphi_k \otimes \phi\|_2 = \|(1 - q_{t_i})\varphi_k\|_2 \|\phi\|_2 = \|(1 - q_{t_i})\varphi_k\|_2 \quad \text{and} \quad \|\varphi_k \otimes \phi\|_2 = \|\varphi_k\|_2 \|\phi\|_2 = 1,$$

so for $1 \leq i, j \leq n$ we have

$$\|\bar{Q}_{t_i} e^{(t_j - t_i)D} K_{\text{GUE}, N}\|_1 \leq \sum_{k=0}^{N-1} e^{(t_j - t_i)k} \|\bar{Q}_{t_i}(\varphi_k \otimes \varphi_k)\|_1 \leq \sum_{k=0}^{n-1} e^{(t_j - t_i)k} \|\bar{q}_{t_i} \varphi_k\|_2 < \infty,$$

since $|\varphi_k(x)| \leq cx^k e^{-x^2/2}$ and $|\bar{q}_{t_i}(x)| \leq ce^{\kappa x^2}$ where $\kappa < \frac{1}{2\sqrt{2}} \tanh(\tau/\sqrt{2}) < \frac{1}{2}$. Hence the only thing left to check in (ii) is that $\|\bar{Q}_{t_i} e^{-(t_j - t_i)D}\|_1 < \infty$ for $i < j$. To that end we use the Feynman-Kac representation to write (setting $t = \frac{1}{2}(t_j - t_i)$)

$$e^{-tD}(x, y) = \frac{1}{\sqrt{2\pi t}} e^{-(x-y)^2/2t} \mathbb{E}_{b(0)=x, b(t)=y} \left[e^{-\frac{1}{2} \int_0^t (b(s)^2 - 1) ds} \right], \quad (52)$$

where $b(s)$ denotes a standard Brownian motion and the subscript in the expectation means that it is conditioned (in the sense of a Brownian bridge) to go from x at time 0 to y at time t . Then

$$\begin{aligned} \|\bar{Q}_{t_i} e^{-tD}\|_2^2 &= \int_{\mathbb{R}^2} dx dy \bar{q}_{t_i}(x)^2 \frac{1}{2\pi t} e^{-(x-y)^2/t} \mathbb{E}_{b(0)=x, b(t)=y} \left[e^{-\frac{1}{2} \int_0^t (b(s)^2 - 1) ds} \right]^2 \\ &\leq \int_{\mathbb{R}^2} dx dy c e^{2\kappa x^2} \frac{1}{2\pi t} e^{-(x-y)^2/2t} \mathbb{E}_{b(0)=x, b(t)=y} \left[e^{-\int_0^t (b(s)^2 - 1) ds} \right] \\ &\leq \int_{-\infty}^{\infty} dx c \frac{e^{2\kappa x^2 + t}}{\sqrt{2\pi t}} \mathbb{E}_{b(0)=x} \left[e^{-\int_0^t b(s)^2 ds} \right] \leq c' \int_{-\infty}^{\infty} dx e^{2\kappa x^2 - \tanh(\sqrt{2}t)x^2/\sqrt{2}} < \infty \end{aligned}$$

by our assumption on κ , where in the last inequality we have used (1.9.3) in [18]. In the same way we have $\|e^{-tD}\|_2 < \infty$ and then $\|\bar{Q}_{t_i} e^{-(t_j - t_i)D}\|_1 \leq \|\bar{Q}_{t_i} e^{-tD}\|_2 \|e^{-tD}\|_2 < \infty$.

Finally, Assumption 3(iii) follows from Assumption 3(ii) thanks to the observation in Remark 3.2. \square

Proof of Proposition 4.3. Using the notation introduced before the statement of the result, it is clear that the functions q_{t_i} satisfy the assumptions appearing in Corollary 4.2, so that

$$\mathbb{E} \left[\prod_{j=1}^N \prod_{i=1}^n (1 - \delta h(t_j, \lambda_j^N(t_i))) \right] = \det(I - K_{\text{GUE},N} + Q_{t_1} e^{(t_1-t_2)D} \dots Q_{t_n} e^{(t_n-t_1)D} K_{\text{GUE},N})_{L^2(\mathbb{R})}. \quad (53)$$

The left-hand side equals

$$\begin{aligned} \mathbb{E} \left[\prod_{j=1}^N \exp \left(\sum_{i=1}^n [\log(1 - \delta h(t_i, \lambda_j^N(t_i)))] \right) \right] &= \mathbb{E} \left[\prod_{j=1}^N \exp \left(-\delta \sum_{i=1}^n h(t_i, \lambda_j^N(t_i)) + n\mathcal{O}(\delta^2) \right) \right] \\ &\xrightarrow{n \rightarrow \infty} \mathbb{E} \left[\prod_{j=1}^N \exp \left(-\int_{\ell}^r h(t, \lambda_j^N(t)) dt \right) \right] \end{aligned} \quad (54)$$

by the dominated convergence theorem.

For the right-hand side of (53), writing $\Gamma_{\ell,r}^{h,n} = Q_{t_1} e^{(t_1-t_2)D} Q_{t_2} \dots e^{(t_{n-1}-t_n)D} Q_{t_n}$ one can use the Feynman-Kac representation on each interval $[t_i, t_{i+1}]$ as in (52) (see also (3)) to deduce that $\Gamma_{\ell,r}^{h,n}$ has kernel

$$\Gamma_{\ell,r}^{h,n}(x, y) = \frac{1}{\sqrt{2\pi(r-\ell)}} e^{-(x-y)^2/2(r-\ell)} \mathbb{E}_{b(\ell)=x, b(r)=y} \left[e^{\sum_{i=0}^n \log(1-\delta h(t_i, b(t_i))) - \frac{1}{2} \int_{\ell}^r (b(s)^2 - 1) ds} \right],$$

where $b(s)$ is a Brownian bridge (with diffusion coefficient 2) run from x at time ℓ to y at time r . Then using (36) we deduce that

$$\begin{aligned} [\Gamma_{\ell,r}^{h,n} - \Gamma_{\ell,r}^h](x, y) &= \frac{1}{\sqrt{2\pi(r-\ell)}} e^{-(x-y)^2/2(r-\ell)} \mathbb{E}_{b(r)=x, b(\ell)=y} \left[e^{-\frac{1}{2} \int_{\ell}^r (2h(s, b(s)) + b(s)^2 - 1) ds} \right. \\ &\quad \left. \cdot \left(e^{\sum_{i=0}^n \log(1-\delta h(t_i, b(t_i))) + \int_{\ell}^r h(s, b(s)) ds} - 1 \right) \right] \end{aligned}$$

for small enough δ . Since h is bounded and continuous, the random variable inside the expectation goes to 0 almost surely as $n \rightarrow \infty$ using a similar argument as in (54), and thus since this random variable is bounded by $ce^{-\int_{\ell}^r b(s)^2 ds}$ the whole expected value goes to 0 as $n \rightarrow \infty$ by the dominated convergence theorem. If we now define the multiplication operator $Mf(x) = \phi(x)f(x)$ with $\phi(x) = (1+x^2)^{-1/2}$ then the above argument gives $(\Gamma_{\ell,r}^{h,n} - \Gamma_{\ell,r}^h)M(x, y) \rightarrow 0$ as $n \rightarrow \infty$ for all x, y . To deduce that $\|(\Gamma_{\ell,r}^{h,n} - \Gamma_{\ell,r}^h)M\|_2 \rightarrow 0$ as $n \rightarrow \infty$ we use the dominated convergence theorem again together with the fact that $(\Gamma_{\ell,r}^{h,n} - \Gamma_{\ell,r}^h)M$ satisfies

$$\begin{aligned} \int_{\mathbb{R}^2} dx dy [(\Gamma_{\ell,r}^{h,n} - \Gamma_{\ell,r}^h)M(x, y)]^2 &\leq c \int_{\mathbb{R}^2} dx dy e^{-\frac{(x-y)^2}{r-\ell}} \mathbb{E}_{b(\ell)=x, b(r)=y} \left[e^{-\frac{1}{2} \int_{\ell}^r b(s)^2 ds} \right]^2 \phi(y)^2 \\ &\leq c \left[\int_{-\infty}^{\infty} dy \phi(y)^4 \right]^{1/2} \int_{-\infty}^{\infty} dx \left[\int_{-\infty}^{\infty} dy e^{-\frac{2(x-y)^2}{r-\ell}} \mathbb{E}_{b(\ell)=x, b(r)=y} \left[e^{-2 \int_{\ell}^r b(s)^2 ds} \right] \right]^{1/2} \\ &\leq c \|\phi\|_4^2 \int_{-\infty}^{\infty} dx \left[\int_{-\infty}^{\infty} dy e^{-\frac{(x-y)^2}{2(r-\ell)}} \mathbb{E}_{b(\ell)=x, b(r)=y} \left[e^{-2 \int_{\ell}^r b(s)^2 ds} \right] \right]^{1/2} \\ &\leq c \|\phi\|_4^2 \int_{-\infty}^{\infty} dx \mathbb{E}_{b(\ell)=x} \left[e^{-2 \int_{\ell}^r b(s)^2 ds} \right]^{1/2} = c' \|\phi\|_4^2 \left[\int_{-\infty}^{\infty} dx e^{-\tanh(2(r-\ell))x^2} \right]^{1/2} < \infty, \end{aligned}$$

where we have used the Cauchy-Schwartz inequality and the last equality follows from (1.9.3) of [18]. Checking that $\|M^{-1}e^{(r-\ell)D}K_{\text{GUE},N}\|_2 < \infty$ is simple as in the proof of

Corollary 4.2, so from the above we deduce that

$$\begin{aligned} & \left\| (K_{\text{GUE},N} - \Gamma_{\ell,r}^{h,n} e^{(r-\ell)D} K_{\text{GUE},N}) - (K_{\text{GUE},N} - \Gamma_{\ell,r}^h e^{(r-\ell)D} K_{\text{GUE},N}) \right\|_1 \\ & \leq \left\| (\Gamma_{\ell,r}^h - \Gamma_{\ell,r}^{h,n}) M \right\|_2 \left\| M^{-1} e^{(r-\ell)D} K_{\text{GUE},N} \right\|_2 \xrightarrow{n \rightarrow \infty} 0. \end{aligned}$$

Since the mapping $A \mapsto \det(I + A)_{L^2(\mathbb{R})}$ is continuous in the space of trace class operators (see [43]), we deduce that the right-hand side of (53) converges to $\det(I - K_{\text{GUE},N} + \Gamma_{\ell,r}^h e^{(r-\ell)D} K_{\text{GUE},N})_{L^2(\mathbb{R})}$, and hence

$$\mathbb{E} \left[\prod_{j=1}^N \exp \left(- \int_{\ell}^r h(t, \lambda_j^N(t)) dt \right) \right] = \det(I - K_{\text{GUE},N} + \Gamma_{\ell,r}^h e^{(t_n - t_1)D} K_{\text{GUE},N})_{L^2(\mathbb{R})}. \quad \square$$

Proof of Corollary 4.6. Fix $f \in L^2(\mathbb{R})$ and write $\phi(x) = e^{rx} \mathbf{1}_{x \geq 0} + \varphi(x)^{-1} \mathbf{1}_{x < 0}$. Then for $i < j$ (note that as in the proof of Corollary 4.2 we need to consider the operators with bars), writing $\hat{f}(\lambda) = \int_{-\infty}^{\infty} dx \text{Ai}(x + \lambda) f(x)$ we have

$$\begin{aligned} \|\overline{Q}_{t_i} e^{(t_i - t_j)H} f\|_2^2 & \leq c \int_{-\infty}^{\infty} dx \left[\int_{\mathbb{R}^2} dy d\lambda \phi(x) e^{\lambda(t_i - t_j)} \text{Ai}(x + \lambda) \text{Ai}(y + \lambda) f(y) \right]^2 \\ & \leq c \int_{-\infty}^{\infty} dx \left[\int_{-\infty}^{\infty} d\lambda \phi(x)^2 e^{2\lambda(t_j - t_i)} \text{Ai}(x + \lambda)^2 \right] \left[\int_{-\infty}^{\infty} d\lambda \hat{f}(\lambda)^2 \right] \\ & = c \|f\|_2^2 \int_{-\infty}^{\infty} dx \phi(x)^2 e^{-2(t_j - t_i)x} \int_{-\infty}^{\infty} d\lambda e^{2(\lambda+1)(t_j - t_i)} \text{Ai}(\lambda)^2 \leq c' \|f\|_2^2. \end{aligned} \quad (55)$$

where in the last line we have used the Parseval identity for the Airy transform $\int \hat{f}^2 = \int f^2$. The fact that $c' < \infty$ follows from the assumption on r and φ and the bounds

$$|\text{Ai}(x)| \leq c e^{-\frac{2}{3}x^{3/2}} \quad \text{for } x \geq 0 \quad \text{and} \quad |\text{Ai}(x)| \leq c|x|^{-1/4} \quad \text{for } x < 0 \quad (56)$$

(see (10.4.59-60) in [1]). This shows that for $i < j$, $\overline{Q}_{t_i} e^{-(t_j - t_i)H}$ is a bounded operator mapping $L^2(\mathbb{R})$ to itself. Similar computations allow to check the rest of Assumption 1(i). To check (ii) we use the formula given in Lemma 3.1. Each term can be written as a product of the form $(-1)^k (e^{(t_1 - t_{a_0})H} \overline{Q}_{t_{a_0}}) \dots (e^{(t_{a_{k-1}} - t_{a_k})H} \overline{Q}_{t_{a_k}}) (e^{(t_{a_k} - t_{a_1})H} K_2)$ with $1 \leq a_0 < \dots < a_k \leq n$. The $k+1$ factors coming after $(-1)^k$ can be checked to be bounded operators on $L^2(\mathbb{R})$ by a computation similar to (55), and a simpler computation gives the same for $e^{(t_{a_k} - t_{a_1})H} K_2$ using the spectral formula for its kernel.

As in the previous example, checking Assumption 2 is straightforward. For Assumption 3 we choose

$$V_{t_i} f(x) = U_{t_i}' f(x) = \psi(x) f(x), \quad V_{t_i}' f(x) = U_{t_i} f(x) = \psi(x)^{-1} f(x)$$

with $\psi(x) = e^{-rx/2} \mathbf{1}_{x \geq 0} + \varphi(x)^{1/2} \mathbf{1}_{x < 0}$. Condition (i) is obvious. Now note that $K_2 = B_0 P_0 B_0$, where $B_0(x, \lambda) = \text{Ai}(x + \lambda)$ and $P_a f(x) = f(x) \mathbf{1}_{x \geq a}$, so

$$\|V_{t_i} \overline{Q}_{t_i} K_2 V_{t_i}'\|_1 \leq \|V_{t_i} \overline{Q}_{t_i} B_0 P_0\|_2 \|P_0 B_0 V_{t_i}'\|_2. \quad (57)$$

We have

$$\begin{aligned} \|V_{t_i} \overline{Q}_{t_i} B_0 P_0\|_2^2 & \leq c \int_{-\infty}^{\infty} dx \int_0^{\infty} d\lambda \psi(x)^2 \phi(x)^2 \text{Ai}(x + \lambda)^2 \\ & = c \int_0^{\infty} dx \int_0^{\infty} d\lambda e^{rx} \text{Ai}(x + \lambda)^2 + c \int_{-\infty}^0 dx \varphi(x)^{-1} \int_{-\infty}^{\infty} d\lambda \text{Ai}(\lambda)^2. \end{aligned}$$

The first integral on the right-hand side is clearly finite by (56). For the second one, note that by (56) $\int_{-\infty}^{\infty} d\lambda \text{Ai}(\lambda)^2 \leq c(1 + |x|^{1/2})$, so the integral is finite by the assumption on φ ,

which by the Cauchy-Schwartz inequality implies $\int_{-\infty}^0 dx |x|^{1/2} \varphi(x)^{-1} < \infty$. $\|P_0 B_0 V'_{t_i}\|_2 < \infty$ follows from the exact same calculation, and hence from (57) we get that $V_{t_i} \overline{Q}_{t_i} K_2 V'_{t_i}$ is trace class. The same proof shows that $V_{t_i} \overline{Q}_{t_i} e^{(t_j - t_i)H} K_2 V'_{t_i}$ is trace class for $i < j$. To check that $V_{t_i} \overline{Q}_{t_i} e^{-tH} V'_{t_j}$ is trace class for $i < j$ and $t = t_j - t_i > 0$ we start by writing

$$\|V_{t_i} \overline{Q}_{t_i} e^{-tH} V'_{t_j}\|_2 \leq \|V_{t_i} \overline{Q}_{t_i} e^{-tH/2}\|_2 \|e^{-tH/2} V'_{t_j}\|_2.$$

For the first factor we use again the explicit formula for the kernel of e^{-tH} to obtain

$$\begin{aligned} \|V_{t_i} \overline{Q}_{t_i} e^{-tH/2}\|_2^2 &= \int_{\mathbb{R}^4} dx dy d\lambda d\sigma \psi(x)^2 \phi(x)^2 e^{t(\lambda+\sigma)/2} \text{Ai}(x+\lambda) \text{Ai}(y+\lambda) \text{Ai}(x+\sigma) \text{Ai}(y+\sigma) \\ &= \int_{-\infty}^{\infty} dx \int_{-\infty}^{\infty} d\lambda \psi(x)^2 \phi(x)^2 e^{t\lambda} \text{Ai}(x+\lambda)^2 = \int_{-\infty}^{\infty} dx \psi(x)^2 \phi(x)^2 e^{-tx} \int_{-\infty}^{\infty} d\lambda e^{t\lambda} \text{Ai}(\lambda)^2 \end{aligned} \quad (58)$$

which is finite by the similar arguments as above. $\|e^{-tH/2} V'_{t_j}\|_2$ can be bounded in the same manner, and we deduce that $V_{t_i} \overline{Q}_{t_i} e^{-tH} V'_{t_j}$ is trace class. The same proof works for $V_{t_i} \overline{Q}_{t_i} e^{-tH} K_2 V'_{t_j}$.

To get (iii) we use Lemma 3.1 and rewrite each term in the sum as

$$\begin{aligned} (-1)^k (U_{t_i} e^{(t_i - t_{a_0})H} \overline{Q}_{t_{a_0}} U'_{t_{a_0}}) (U_{t_{a_0}} e^{(t_{a_0} - t_{a_1})H} \overline{Q}_{t_{a_1}} U'_{t_{a_1}}) \cdots (U'_{t_{a_{k-1}}} e^{(t_{a_{k-1}} - t_{a_k})H} \overline{Q}_{t_{a_k}} U'_{t_{a_k}}) \\ \cdot (U_{t_{a_k}} e^{(t_{a_k} - t_{a_1})H} K_2 U'_{t_{a_1}}). \end{aligned}$$

Since $U_{t_i} = V'_{t_i}$ and $U'_{t_i} = V_{t_i}$, each factor above corresponds to the adjoint of one of the factors appearing in (ii). Since the adjoint of a trace class operator is also trace class, we deduce that the whole product is trace class. \square

Proof of Corollary 4.8. We already indicated how to check Assumption 2. One checks directly that the first three operators in Assumption 1(i) are bounded operators preserving $L^2(\mathbb{R})$, while the last one can be checked using (46) and arguing about the Airy functions appearing there similarly as in the previous proof. Assumption 1(ii) follows similarly using Lemma 3.1. Assumption 3 can be checked following the same ideas as in the proof of Corollary 4.6 and using the arguments in Appendix A of [9] to provide the necessary analytical estimates. \square

REFERENCES

- [1] M. Abramowitz, I. A. Stegun. *Handbook of mathematical functions with formulas, graphs, and mathematical tables*. Vol. 55. National Bureau of Standards Applied Mathematics Series, 1964, pp. xiv+1046
- [2] M. Adler, P.L. Ferrari, P. van Moerbeke. Airy processes with wanderers and new universality classes. *Ann. Probab.*, **38**:714-769 (2010).
- [3] M. Adler, J. Delépine, P. van Moerbeke. Dyson's nonintersecting Brownian motions with a few outliers. *Comm. Pure Appl. Math.*, **62**:334-395 (2009).
- [4] G. W. Anderson, A. Guionnet, O. Zeitouni. *An introduction to random matrices*. Cambridge Studies in Advanced Mathematics, vol. 118, Cambridge University Press, Cambridge (2010).
- [5] A. Aptekarev, P. Bleher, and A. Kuijlaars. Large n limit of Gaussian random matrices with external source, part II. *Comm. Math. Phys.*, **259**:367-389 (2005).
- [6] J. Baik, G. Ben Arous, S. Péché. Phase transition of the largest eigenvalue for nonnull complex sample covariance matrices. *Ann. Probab.*, **33**:1643-1697 (2005).
- [7] A. Borodin. Determinantal point processes. Chapter 11 in *The Oxford Handbook of Random Matrix Theory*, Oxford University Press (2011).
- [8] A. Borodin, M. Duits. Limits of determinantal processes near a tacnode. *Ann. Inst. H. Poincaré B*, **47**:243-258 (2011).

- [9] A. Borodin, P. L. Ferrari, M. Prähofer. Fluctuations in the discrete TASEP with periodic initial configurations and the Airy_1 process. *Int. Math. Res. Pap.*, rpm002, 47 (2007).
- [10] A. Borodin, P. L. Ferrari, M. Prähofer, T. Sasamoto. Fluctuation properties of the TASEP with periodic initial configuration *J. Stat. Phys.* **129**:5-6:1055-1080 (2007).
- [11] A. Borodin, P. L. Ferrari, T. Sasamoto. Transition between Airy_1 and Airy_2 processes and TASEP fluctuations. *Comm. Pure Appl. Math.*, **61**:1603-1629 (2008).
- [12] A. Borodin, G. Olshanski. Point processes and the infinite symmetric group. *Math. Research Lett.*, **5**:799-816 (1998).
- [13] A. Borodin, G. Olshanski. Random partitions and the Gamma kernel. *Adv. Math.*, **194**(1):141-202 (2005).
- [14] A. Borodin, G. Olshanski. Stochastic dynamics related to Plancherel measures on partitions. *Representation Theory, Dynamical Systems, and Asymptotic Combinatorics*. AMS Translations - Series 2: Advances in the Mathematical Sciences, **217**:9-21 (2006).
- [15] A. Borodin, G. Olshanski. Markov processes on partitions. *Probab. Theory Relat. Fields*, **135**:84-152 (2006).
- [16] A. Borodin, S. Péché. Airy kernel with two sets of parameters in directed percolation and random matrix theory. *J. Stat. Phys.*, **132**:275-290 (2008).
- [17] A. Borodin, E. M. Rains. Eynard-Mehta theorem, Schur process, and their Pfaffian analogs. *J. Stat. Phys.*, **121**:291-317 (2005).
- [18] A. N. Borodin, P. Salminen. *Handbook of Brownian Motion: facts and formulae*. Birkhäuser, 2002, 2nd Ed.
- [19] E. Brézin and S. Hikami,. Universal singularity at the closure of a gap in a random matrix theory. *Phys. Rev. E*, **57**:4140-4149 (1998).
- [20] E. Brézin and S. Hikami. Level Spacing of Random Matrices in an External Source. *Phys. Rev. E* , **58**:7176-7185 (1998).
- [21] I. Corwin, A. Hammond. Brownian Gibbs property for Airy line ensembles. arXiv:1108.2291.
- [22] I. Corwin, J. Quastel, D. Remenik. Continuum statistics of the Airy_2 process. *Comm. Math. Phys.*, **317**:347-362 (2013).
- [23] B. Eynard, M. L. Mehta. Matrices coupled in a chain. I. Eigenvalue correlations. *J. Phys. A: Math. Gen.*, **31**:4449-4456 (1998).
- [24] P. L. Ferrari. The universal Airy_1 and Airy_2 processes in the totally asymmetric simple exclusion process. *Integrable Systems and Random Matrices: In Honor of Percy Deift*, Contemporary Mathematics **458**:321-332 (2008).
- [25] I. Gessel, G. Viennot. Binomial determinants, paths, and hook length formulae. *Adv. in Math.* **58**:300-321 (1985).
- [26] K. Johansson. Discrete polynuclear growth and determinantal processes. *Comm. Math. Phys.*, **242**:277-329 (2003).
- [27] K. Johansson. Non-intersecting paths, random tilings and random matrices. *Probab. Theory Relat. Fields*, **123**:225-280 (2002).
- [28] I. Karatzas, S. Shreve. *Brownian motion and stochastic calculus*. Springer, 1991.
- [29] S. Karlin, G. McGregor. Coincidence probabilities. *Pacific J. Math.* **9**:1141-1164 (1959).
- [30] M. Katori, H. Tanemura. Noncolliding squared Bessel processes. *J. Stat. Phys.*, **142**:592-615 (2011).
- [31] B. Lindström. On the vector representation of induced matroids. *Bull. London Math. Soc.* **5**:85-90 (1973).
- [32] G. Moreno Flores, J. Quastel, D. Remenik. Endpoint distribution of directed polymers in 1+1 dimensions. *Comm. Math. Phys.*, **317**:363-380 (2013).
- [33] T. Nagao, P. J. Forrester. Multilevel dynamical correlation function for Dyson's Brownian motion model of random matrices. *Phys. Lett. A*, **247**:42-46 (1998).
- [34] A. Okounkov and N. Reshetikhin. Correlation function of Schur process with application to local geometry of a random 3-dimensional Young diagram. *J. Amer. Math. Soc.*, **16**:581-603 (2003).
- [35] G. Olshanski. An introduction to harmonic analysis on the infinite symmetric group. *Asymptotic Combinatorics with Applications to Mathematical Physics (A. Vershik, ed.)*. Springer Lecture Notes in Math. **1815**:127-160, (2003).
- [36] M. Prähofer, H. Spohn. Scale invariance of the PNG droplet and the Airy process. *J. Stat. Phys.*, **108**:1071-1106 (2002).
- [37] S. Prohac, H. Spohn. The one-dimensional KPZ equation and the Airy process. *J. Stat. Mech. Theor. Exp.*, **03**:P03020 (2011).
- [38] J. Quastel, D. Remenik. Tails of the endpoint distribution of directed polymers. *Ann. Inst. Henri Poincaré Probab. Stat.*, to appear.

- [39] J. Quastel, D. Remenik. Supremum of the Airy_2 process minus a parabola on a half-line. *J. Stat. Phys.*, <http://dx.doi.org/10.1007/s10955-012-0633-4>.
- [40] J. Quastel, D. Remenik. Local behavior and hitting probabilities of the Airy_1 process. *Probab. Theory Relat. Fields*, <http://dx.doi.org/10.1007/s00440-012-0466-8> (2012).
- [41] J. Quastel, D. Remenik. Airy processes and variational problems. arXiv:1301.0750.
- [42] T. Sasamoto. Spatial correlations of the 1D KPZ surface on a flat substrate. *J. Phys. A: Math. Gen.*, **38**:L549 (2005)
- [43] B. Simon. *Trace Ideals and Their Applications*. AMS, 2nd ed. (2000).
- [44] J. R. Stembridge. Nonintersecting paths, pfaffians, and plane partitions. *Adv. in Math.*, **83**:96-131 (1990).
- [45] C. Tracy, H. Widom. Differential equations for the Dyson process. *Comm. Math. Phys.*, **252**:7-41 (2004).
- [46] C. Tracy, H. Widom. The Pearcey process. *Comm. Math. Phys.*, **263**:381-400 (2006).

A. BORODIN, MASSACHUSETTS INSTITUTE OF TECHNOLOGY, DEPARTMENT OF MATHEMATICS, 77 MASSACHUSETTS AVENUE, CAMBRIDGE, MA 02139-4307, USA AND INSTITUTE FOR INFORMATION TRANSMISSION PROBLEMS, MOSCOW, RUSSIA

E-mail address: `borodin@math.mit.edu`

I. CORWIN, MASSACHUSETTS INSTITUTE OF TECHNOLOGY, DEPARTMENT OF MATHEMATICS, 77 MASSACHUSETTS AVENUE, CAMBRIDGE, MA 02139-4307, USA

E-mail address: `ivan.corwin@gmail.com`

D. REMENIK, DEPARTAMENTO DE INGENIERÍA MATEMÁTICA AND CENTRO DE MODELAMIENTO MATEMÁTICO, UNIVERSIDAD DE CHILE, AV. BLANCO ENCALADA 2120, SANTIAGO, CHILE

E-mail address: `dremenik@dim.uchile.cl`